

## PRODUCT INFORMATION

# OGIVRI

*Trastuzumab (rch)*



## NAME OF THE MEDICINE

Active ingredient:	trastuzumab
Chemical name:	Immunoglobulin G1, anti-human p185 neu receptor human-mouse monoclonal recombinant human monoclonal antibody (r-hu-MAb) human epidermal growth factor receptor 2 (HER2) g1-chain, disulfide with human-mouse monoclonal r-hu-MAb HER2 light chain, dimer
Molecular formula:	$C_{6460}H_{9972}N_{1724}O_{2014}S_{44}$
Molecular weight:	~148 kDa
CAS Registry Number:	180288-69-1

## DESCRIPTION

OGIVRI (trastuzumab) is a recombinant DNA-derived humanized monoclonal antibody that selectively targets the extracellular domain of the human epidermal growth factor receptor 2 protein (HER2). The antibody is an IgG<sub>1</sub> kappa that contains human framework regions with the complementarity-determining regions of a murine anti-p185 HER2 antibody that binds to HER2. Trastuzumab is composed of 1,328 amino acids and has a molecular weight of ~148 kDa.

The humanized antibody against HER2 is produced by recombinant mammalian cells (Chinese hamster ovary (rch)) in suspension culture in a nutrient medium and purified by affinity chromatography and ion exchange, including specific viral inactivation and removal procedures.

OGIVRI is a sterile, white to pale yellow, preservative-free lyophilized powder for IV infusion.

OGIVRI is available as a single-dose vial containing 150 mg of trastuzumab or as a Pharmacy Bulk Pack, containing 440 mg of trastuzumab with the following excipients: L-histidine hydrochloride monohydrate, L-histidine, sorbitol and Macrogol 3350.

OGIVRI is a biosimilar medicine to Herceptin.

### Excipient with Known Effect

#### Sorbitol

OGIVRI contains 0.77 mg sorbitol/ 1.00 mg trastuzumab. There is 115.2 mg of sorbitol in the 150 mg vial and 337.9 mg of sorbitol in the 440 mg Pharmacy Bulk Pack

Reconstitution of the 150 mg vial with 7.2 mL of sterile water for injection yields 7.4 mL of a single-dose solution containing approximately 21 mg/mL trastuzumab, at a pH of approximately 6.0. A volume overage of 4% ensures that the labelled dose can be withdrawn from each vial.

Reconstitution of the 440 mg vial with 20 mL of bacteriostatic water for injection yields a multiple-dose solution containing approximately 21 mg/mL trastuzumab, at a pH of approximately 6.0.

## PHARMACOLOGY

### Pharmacodynamics

The HER2 (or c-erbB2) proto-oncogene encodes for a single transmembrane spanning, receptor-like protein of 185 kDa, which is structurally related to the epidermal growth factor receptor.

Trastuzumab has been shown, both in *in-vitro* assays and in animals, to inhibit the proliferation of human tumour cells that overexpress HER2. *In vitro*, trastuzumab-mediated antibody-dependent cell-mediated cytotoxicity (ADCC) has been shown to be preferentially exerted on HER2 overexpressing cancer cells compared with cancer cells that do not overexpress HER2. In animal models *in vivo*, murine anti-HER2 antibody inhibited the growth of human tumours overexpressing HER2, indicating that the humanized antibody (trastuzumab) is likely also to have anti-proliferative activity *in vivo* against human breast tumours expressing high levels of HER2.

### Pharmacokinetics

The pharmacokinetics of trastuzumab have been studied in patients with breast cancer (metastatic and early) and advanced gastric cancer (AGC).

The pharmacokinetics of trastuzumab were evaluated in a population pharmacokinetic model analysis using pooled data from 1,582 from 18 Phase I, II and III trials receiving trastuzumab IV to treat a range of cancers, but mostly breast and gastric cancer. A two-compartment model with parallel linear and non-linear elimination from the central compartment described the trastuzumab concentration-time profile. Due to the non-linear elimination, total clearance increased with decreasing concentrations. Linear clearance was 0.127 L/day for breast cancer (metastatic and early) and 0.176 L/day for AGC. The nonlinear elimination parameter values were 8.81 mg/day for the maximum elimination rate ( $V_{max}$ ) and 8.92 mg/L for the Michaelis-Menten constant ( $K_m$ ). The central compartment volume was 2.62 L for patients with breast cancer and 3.63 L for patients with AGC.

The population predicted PK exposures (with 5<sup>th</sup> - 95<sup>th</sup> Percentiles) and PK parameter values at clinically relevant concentrations ( $C_{max}$  and  $C_{min}$ ) for breast cancer and AGC patients treated with the approved q1w and q3w dosing regimens are shown in Table 1 (Cycle 1) and Table 2 (steady-state) below.

**Table 1: Population Predicted Cycle 1 PK Exposure Values (with 5<sup>th</sup> - 95<sup>th</sup> Percentiles) for IV Regimens in Breast Cancer and AGC Patients**

Regimen	Primary tumor type	N	Cmin
---------	--------------------	---	------

Regimen	Primary tumor type	N	Cmin,ss	Cmax,ss	AUCss	Time to steady-state (week)	Total CL range at steady-state (L/day)
8mg/kg + 6mg/kg q3w	MBC/EBC	1195	47.4 (5.0 - 114.7)	179.4 (107.3-308.8)	1794.2 (673.0 – 3618.4)	12	0.173 - 0.283
	AGC	274	32.9 (6.1 - 88.9)	131.0 (72.5 - 250.5)	1338.2 (557.0- 2875.4)	9	0.189 - 0.337
4mg/kg + 2mg/kg qw	MBC/EBC	1195	66.1 (14.9-142.3)	108.8 (51.0 - 208.6)	1765.3 (647.3 – 3578.1)	12	0.201 - 0.244

### Pharmacokinetics in Special Populations

Dedicated pharmacokinetic studies in the elderly and those with renal or hepatic impairment have not been carried out. However, in a population PK analysis, age and renal impairment were not shown to affect trastuzumab disposition. The population PK analysis showed that the estimated creatinine clearance (Cockcroft and Gault) does not correlate with the pharmacokinetics of trastuzumab.

## CLINICAL TRIALS

### Early Breast Cancer

Early breast cancer is defined as non-metastatic, primary, invasive carcinoma of the breast.

### Trastuzumab in Combination with Adjuvant Chemotherapy

The use of trastuzumab in the setting of early breast cancer (after surgery and in association with chemotherapy and, if applicable, radiotherapy) has been studied in four multicentre randomized phase III trials of patients with HER2 positive breast cancer who have completed surgery. In these clinical trials, early breast cancer was limited to operable, primary adenocarcinoma of the breast with positive axillary nodes or node negative disease with additional indicators of a higher degree of risk. The design of these studies is summarized in Table 3 and efficacy results are presented in Tables 4-8.

**Table 3: Clinical Trials in Early Breast Cancer**

	HERA trial n = 3386	NSAPB B-31 and NCCTG N9831 trials (joint analysis) n = 3763	BCIRG 006 n = 3222
<b>Eligible patients</b>	Node positive or node negative [n = 1098] and tumour size >1 cm;  <i>Protocol initially unrestricted but amended and node negative patients with 8.5%] and node negative patients with tumours &gt;1 509,46.4%] were included</i>	Node positive or node negative [n = 190] and tumour size  >2 cm regardless of hormonal status; or >1 cm and ER–ve  [n = 63 node-negative	Node positive or node negative and at least 1 of the following: tumour size > 2 cm and ER and PR -ve, or histologic and/or nuclear grade 2-3, or age < 35 years.

OGIVRI – PRODUCT INFORMATION

4

	<b>HERA trial n = 3386</b>	<b>NSAPB B-31 and NCCTG N9831 trials (joint analysis) n = 3763</b>	<b>BCIRG 006 n = 3222</b>
<b>Trastuzumab dosage regimen</b>	Loading dose 8 mg/kg, followed by 6 mg/kg (q3w)	Loading dose 4 mg/kg, followed by 2 mg/kg (q1w)	Loading dose 4 mg/kg, followed by 2 mg/kg (q1w). After chemo, 6 mg/kg (q3w)
<b>Duration of trastuzumab treatment</b>	1 yr or 2 yrs	52 weeks	52 weeks
<b>Chemotherapy regimen(s)</b>	Various HERA trial n = 3386	AC (q3w) followed by IV paclitaxel as a NSAPB B-31 and NCCTG N9831 trials (joint analysis) n = 3763	AC followed by BCIRG 006 n = 3222
		continuous IV infusion  Paclitaxel: 80 mg/m <sup>2</sup> q1w for 12 weeks or 175 mg/m <sup>2</sup> q3w for 4 cycles (day 1 of each cycle)	docetaxel and carboplatin (DCarb)  Docetaxel (IV infusion over 60 min): q3w for 4 cycles or (DCarb): 75 mg/m <sup>2</sup> q3w for 6 cycles Carboplatin (at target AUC): 6 mg/mL/min (IV infusion over 30 - 60 min) q3w for a total of 6 cycles.
<b>Timing of trastuzumab in relation to chemotherapy</b>	After completion of (neo)adjuvant <sup>a</sup>		and DCarbH)
<b>Median follow-up</b>	1 year (initial evaluation) [8 years (follow-up evaluation)]	2 years	3 years

AC = doxorubicin + cyclophosphamide; q3w = every 3 weeks; q1w = weekly chemo = chemotherapy; <sup>a</sup> 89% of subjects received adjuvant chemotherapy; 5% received neoadjuvant chemotherapy and 6% received a combination of neoadjuvant and adjuvant chemotherapy.

The HERA trial was designed to compare 1 and 2 years of 3-weekly trastuzumab treatment vs. observation in patients with HER2 positive breast cancer following surgery, established chemotherapy and radiotherapy (if applicable). In addition, a comparison of 2 years trastuzumab treatment vs. 1 year trastuzumab treatment was performed. Patients assigned to receive trastuzumab were given an initial loading dose of 8 mg/kg, followed by 6 mg/kg every 3 weeks for either 1 or 2 years. The efficacy results from the HERA trial are summarized in the following table:

**Table 4: Efficacy Results from the HERA Trial at 12 months<sup>1</sup> and 8 years<sup>2</sup> of median follow up**

Parameter	Observation	Trastuzumab 1yr treatment	p-value	HR (95% CI)
<b>Disease free survival</b> No. of patients with event (1 year <sup>1</sup> )	12.9%	7.5%	<0.0001	0.54 (0.44, 0.67)

No. of patients with event (8 year <sup>2</sup> )	33.6%	27.7%	<0.0001	0.76 (0.67, 0.86)
<b>Overall Survival</b>				
No. of patients with event (1 year <sup>1</sup> )	2.4%	1.8%	0.24	0.75 (0.47, 1.21)
No. of patients with event (8 year <sup>2</sup> )	20.6%	16.3%	0.0005	0.76 (0.65, 0.88)

HR: Hazard ratio; <sup>1</sup> co-primary endpoint of DFS of 1 year vs. observation met the pre-defined statistical boundary; <sup>2</sup> final analysis (includes crossover of 52% of patients from the observation arm to trastuzumab)

The HERA trial included a subgroup of patients ( $n = 602$ ) with small tumours (<2 cm) and node-negative disease. In this subgroup, the relative risk reduction was similar to the overall trial population (HR = 0.50; 95% CI 0.21 - 1.15). However, the benefit in terms of absolute difference in rate of recurrence after 1 year of follow-up was smaller (2.7% recurrence rate with trastuzumab vs. 5.5% with observation).

In the final analysis (8 year median follow up) extending trastuzumab treatment for a duration of 2 years did not show additional benefit over treatment for 1 year [DFS HR in the intent to treat (ITT) population of 2 years vs. 1 year = 0.99 (95% CI: 0.87, 1.13); p-value = 0.90 and OS HR = 0.98 (0.83, 1.15); p-value = 0.78]. The rate of asymptomatic cardiac dysfunction was increased in the 2-year treatment arm (8.1% vs. 4.6% in the 1-year treatment arm). More patients experienced at least one grade 3 or 4 adverse event in the 2-year treatment arm (20.4%) compared with the 1-year treatment arm (16.3%).

The efficacy results from the joint analysis of the NCCTG 9831 and NSABP B-31 trials are summarized in the following tables:

**Table 5: Summary of Efficacy Results from NSABP B-31 and NCCTG N9831 trials (joint analysis) at the time of the definitive DFS analysis\***

Parameter			p-value	HR (95% CI)
<b>Disease recurrence</b>				
Rate (trastuzumab vs. observation)	15.5%	8.0%	< 0.0001	0.48 (0.39, 0.59)
<b>Survival</b>				
Deaths (trastuzumab vs. observation)	5.5%	3.7%	0.014**	0.67 (0.48, 0.92)

A: doxorubicin; C: cyclophosphamide; P: paclitaxel; H:trastuzumab; HR: Hazard ratio

\*\* p value for OS did not cross the pre-

The pre-planned final analysis of OS from the joint analysis of studies NSABP B-31 and NCCTG N9831 was performed when 707 deaths had occurred (median follow-group). At 8 years, the survival rate was estimated to be 86.9% in the AC + P + H arm and 79.4% in the AC + P arm, an absolute benefit of 7.4% (95% CI 4.9%, 10.0%). The final OS results from the joint analysis of studies NSABP B-31 and NCCTG N9831 are summarized in the following table:

**Table 6: Final Overall Survival Analysis from the joint analysis of trials NSABP B-31 and NCCTG N9831**

Parameter	(N=2032)	(N=2031)	p-value versus	Hazard Ratio (95% CI)
Death (OS event): No. patients with event (%)	418 (20.6%)	289 (14.2%)	< 0.0001	0.64 (0.55, 0.74)

A: doxorubicin; C: cyclophosphamide; P: paclitaxel; H: trastuzumab

The efficacy results from the BCIRG 006 are summarized in the following tables:

**Table 7: Overview of Efficacy Analy**

followed by docetaxel plus trastuzumab; CI = confidence interval

Parameter	= 1073	= 1074	p-value	HR (95% CI)
<b>Disease-free survival (DFS)</b> No. patients with event	195	134	<0.0001	0.61 (0.49, 0.77)
<b>Death (OS event)</b> No. patients with event	80	49	0.0024	0.58 (0.40,0.83)

Parameter	n = 1073	DCarbH n = 1075	p-value	HR (95% CI)
<b>Disease-free survival (DFS)</b> No. patients with event	195	145	0.0003	0.67 (0.54, 0.83)
<b>Death (OS event)</b> No. patients with event	80	56	0.00182	0.66 (0.47, 0.93)

= confidence interval

Based on studies to date, the optimal duration of adjuvant trastuzumab therapy is 1 year and may be clarified in further randomized trials. However, extending adjuvant treatment beyond 1 year is not recommended (see DOSAGE AND ADMINISTRATION/Duration of Treatment).

### Switching treatment from Trastuzumab IV to Trastuzumab SC and vice versa

Study MO22982 (PrefHER) investigated switching from Trastuzumab IV to Trastuzumab SC, and vice versa, in patients with HER2 positive EBC, with a primary objective to evaluate patient preference for either Trastuzumab IV infusion or Trastuzumab SC injection. This trial investigated using a 2-arm, cross-over design with patients being randomized to one of two different q3w Trastuzumab treatment sequences (Trastuzumab IV (Cycles 1- Trastuzumab SC (Cycles 5-8), or Trastuzumab SC (Cycles 1- Trastuzumab IV (Cycles 5-8)). Patients participating in this trial could be enrolled at any time as long as there were at least 10 remaining cycles of Trastuzumab in their planned treatment regimen, therefore patients were either naïve to Trastuzumab IV treatment (20.3%) or pre-exposed to Trastuzumab IV (79.7%) as part of ongoing adjuvant treatment for HER2 positive EBC. Overall, switches from Trastuzumab IV to Trastuzumab SC and vice versa were well tolerated. Pre-switch rates (Cycles 1-4) for SAEs, Grade 3 AEs and treatment discontinuations due to AEs were low (<5%) and similar to post-switch rates (Cycles 5-8). No Grade 4 or Grade 5 AEs were reported. The effect of multiple switches back and forth was not investigated (see also ADVERSE EFFECTS).

### Locally Advanced Breast Cancer

Locally advanced breast cancer is defined as the absence of metastatic disease and meeting one or more of the following criteria: inflammatory breast cancer, a primary tumour that extends to the chest wall or skin, tumour > 5 cm with any positive lymph node(s), any tumour with disease in supraclavicular nodes, infraclavicular nodes or internal mammary nodes, any tumour with axillary lymph nodes fixed to one another or other structures.

### Trastuzumab in Combination with Neoadjuvant-Adjuvant Chemotherapy

The use of trastuzumab for the neoadjuvant-adjuvant treatment of locally advanced breast cancer has been studied in Study MO16432 (NOAH), a multicentre randomized trial, designed to investigate the concurrent administration of trastuzumab with neoadjuvant chemotherapy, including both an



anthracycline and a taxane, followed by adjuvant trastuzumab, up to a total treatment duration of 1 year. The trial recruited patients with newly diagnosed locally advanced (Stage III) or inflammatory breast cancer. Patients with HER2+ tumours were randomized to receive either neoadjuvant chemotherapy concurrently with neoadjuvant-adjuvant trastuzumab ( $n = 116$ ), or neoadjuvant chemotherapy alone ( $n = 118$ ).

Trastuzumab was administered concurrently with 10 cycles of neoadjuvant chemotherapy as follows;

Doxorubicin ( $60 \text{ mg/m}^2$ ) and paclitaxel ( $150 \text{ mg/m}^2$ ) in combination with trastuzumab ( $8 \text{ mg/kg}$  loading dose, followed by  $6 \text{ mg/kg}$  maintenance, administered 3-weekly) for 3 cycles, followed by  
 Paclitaxel ( $175 \text{ mg/m}^2$ ) and trastuzumab ( $6 \text{ mg/kg}$ , administered 3-weekly) for 4 cycles, followed by  
 CMF on day 1 and 8 every 4 weeks for 3 cycles, in combination with 4 cycles of OGIVRI ( $6 \text{ mg/kg}$  administered 3-weekly), followed by  
 up to 7 additional cycles of trastuzumab ( $6 \text{ mg/kg}$ , administered 3-weekly) alone to complete 1 year after starting trastuzumab

The primary endpoint for the trial, event-free survival (EFS), was defined as the time from randomization to disease recurrence or progression (local, regional, distant or contralateral), or death of any cause. The efficacy results from NOAH (full analysis population, defined as all patients who were randomized in the trial following the intent-to-treat principle, with the exception of 3 patients whose data could not be evaluated) are summarized in the table below. The median duration of follow-up in the trastuzumab arm was 3.8 years.

**Table 9: Overview of Efficacy Analyses MO16432 (NOAH)**

Parameter	Chemo + Trastuzumab $n = 115$	Chemo only $n = 116$	p-value	HR (95% CI)
<b>Event-free survival (EFS)</b> No. patients with event	46	59	$p = 0.0275$	0.65 (0.44, 0.96)
<b>Total pathological complete response</b> ^ (95% CI)	40% (31.0, 49.6)	20.7% (13.7, 29.2)	$p = 0.0014$	

^ defined as absence of any invasive cancer both in the breast and axillary nodes; HR: hazard ratio

The addition of trastuzumab to neoadjuvant chemotherapy, followed by adjuvant trastuzumab for a total duration of 52 weeks, resulted in a 35% reduction in the risk of disease recurrence/progression. The hazard ratio translates into an absolute benefit, in terms of 3-year event-free survival rate estimates of 13 percentage points (65 % vs. 52 %) in favour of the trastuzumab arm.

To date, results are not available comparing the efficacy of trastuzumab administered with chemotherapy in the adjuvant setting with that obtained in the neoadjuvant/adjuvant setting.

## Metastatic Breast Cancer

There are no data available to establish the efficacy of trastuzumab for the treatment of metastatic disease in patients who have previously received the medicine for the treatment of early disease.

The safety and efficacy of trastuzumab has been studied in randomized, controlled clinical trials in combination with chemotherapy (Studies H0648g, M77001 and TaNDEM) and in an open-label monotherapy clinical trial (Study H0649g) for the treatment of metastatic breast cancer. All trials studied patients with metastatic breast cancer whose tumours overexpress HER2. Patients were eligible if they had 2+ or 3+ levels of overexpression based on a 0 - 3+ scale by immunohistochemical (IHC) assessment of tumour tissue or whose tumours have HER2 gene amplification as determined by

Fluorescence In Situ Hybridization (FISH) test (see DOSAGE AND ADMINISTRATION, Detection of HER2 Overexpression or HER2 Gene Amplification).

### Trastuzumab in Combination with Chemotherapy


Study H0648g was an open-label, randomized controlled, multinational trial of chemotherapy-alone and in combination with trastuzumab. Patients with previously untreated metastatic breast cancer were treated with either an anthracycline (doxorubicin 60 mg/m<sup>2</sup> or epirubicin 75 mg/m<sup>2</sup>) plus cyclophosphamide (600 mg/m<sup>2</sup>) with or without trastuzumab or paclitaxel (175 mg/m<sup>2</sup> infused over 3 hours) with or without trastuzumab. Patients on trastuzumab treatment received 4 mg/kg intravenous loading dose on Day 0, followed by weekly infusions of 2 mg/kg from Day 7, which they could continue to receive until evidence of disease progression. Patients who had previously received anthracycline based adjuvant therapy were treated with paclitaxel whereas those who were anthracycline naïve were treated with an anthracycline + cyclophosphamide.

The prospectively defined, primary intent-to-treat analysis indicated that the combination of trastuzumab and chemotherapy significantly prolonged time to disease progression (progression-free survival) compared with chemotherapy-alone as first-line treatment of women with metastatic breast cancer who had tumours that overexpressed HER2. The addition of trastuzumab to chemotherapy extended the median time to disease progression by 2.8 months representing a 61% increase ( $p=0.0001$ ).

Both anthracycline-treated and paclitaxel-treated patients benefited from trastuzumab treatment, although the effect appeared to be greater in the paclitaxel stratum. The efficacy of trastuzumab treatment was further supported by the secondary endpoints of response rate, duration of response and one-year survival (see Table 10 below).


One-year survival rates (the prospectively defined survival endpoint) were significantly better for the trastuzumab + chemotherapy versus chemotherapy-alone (79% vs. 68%;  $p=0.008$ ). With a median follow-up of approximately two years, overall survival is improved for patients initially treated with trastuzumab + chemotherapy compared with those receiving chemotherapy-alone (25.4 vs. 20.3 months;  $p=0.025$ ) with a relative risk of death of 0.769 (95% CI 0.607 - 0.973;  $p=0.028$ ).

**Figure 1 Survival Time: Anthracycline ± HERCEPTIN (Study H0648g)**






**Figure 2 Survival Time: Paclitaxel ± HERCEPTIN (Study H0648g)**



The relative overall survival advantage with the addition of trastuzumab was observed in both subgroups: AC [26.8 months (H + AC) vs. 22.8 months (AC-alone);  $p=0.052$ ] and paclitaxel [22.1 months (H + P) vs. 18.4 months (P-alone);  $p=0.273$ ] (see also Figures 1 and 2). The analysis of overall survival was, however, greatly confounded by subsequent trastuzumab treatment of each of control arms' patients, following disease progression, in the open-label extension study, H0659g (59% of patients in the AC-alone group, and 75% of patients in the paclitaxel-alone group subsequently received trastuzumab). Hence, the survival advantage seen above, for trastuzuman + chemotherapy treatment versus chemotherapy-alone (which includes patients who subsequently received trastuzumab) may underestimate the benefit to patients.

Importantly, the efficacy described above was obtained without a significant negative impact on the quality of life. Global quality of life decreased equally in both the chemotherapy-alone group and the trastuzumab + chemotherapy group and was most likely related to the effects of cytotoxic chemotherapy. However, at weeks 20 and 32, the global quality of life score had returned to baseline or better than baseline in the group receiving trastuzumab + chemotherapy, while it remained low in the chemotherapy-alone arm (see Figure 3 below).


**Figure 3 Changes from Baseline in Health-Related Quality-of-Life Scores in Study H0648g**



Study M77001 was a multinational, multi-centre, randomized, controlled trial investigating the safety and efficacy of trastuzumab in combination with docetaxel, as first-line treatment in HER2 positive metastatic breast cancer patients. One hundred and eighty six patients received docetaxel (100 mg/m<sup>2</sup> infused over 1 hour on Day 2) with or without trastuzumab (4 mg/kg loading dose, followed by 2 mg/kg weekly). Sixty percent of patients had received prior anthracycline based adjuvant chemotherapy. trastuzumab with docetaxel was shown to be efficacious in patients whether or not they had received prior adjuvant anthracyclines and regardless of their oestrogen and/or progesterone receptor status.

The combination of trastuzumab + docetaxel significantly increased response rate (61% vs. 34%) and prolonged the median time to disease progression by 4.9 months compared with patients treated with docetaxel-alone (see Table10). Median survival was also significantly increased in patients receiving the combination therapy compared with those receiving docetaxel-alone (30.5 months vs. 22.1 months) (see Figure 4).

**Figure 4 Survival Time: Docetaxel ± trastuzumab (Study M77001)**



**Table 10: Efficacy Outcomes with Combination Therapy for Metastatic Breast Cancer**

	H0648g						M77001	
	H + chemo <i>n</i> = 235	Chemo alone <i>n</i> = 234	H + AC <i>n</i> = 143	AC alone <i>n</i> = 138	H + P <i>n</i> = 92	P alone <i>n</i> = 96	H + D <i>n</i> = 92	D alone <i>n</i> = 94
Median Time to Disease Progression (months, 95% CI)	7.4 (7.0, 9.0)	4.6 (4.4, 5.4)	7.8 (7.3, 9.4)	6.1 (4.9, 7.1)	6.9 (5.3, 9.9)	3.0 (2.1, 4.3)	10.6 (7.6, 12.9)	5.7 (5, 6.5)
<i>p</i> -value <sup>a</sup>	<i>p</i> =0.0001		<i>p</i> =0.0004		<i>p</i> =0.0001		<i>p</i> =0.0001	
Response Rate (%)	50	32	56	42	41	17	61	34
<i>p</i> -value <sup>b</sup>	<i>p</i> <0.0001		<i>p</i> =0.0197		<i>p</i> =0.0002		<i>p</i> =0.0002	
Median Duration of Response (months, 95% CI)	9.1 (7.7,11)	6.1 (5.5,7.8)	9.1 (7.4,12.2)	6.7 (5.8, 8.2)	10.5 (7.3, 12.5)	4.5 (3.9, 6.4)	11.4 (8.3, 15.0)	5.5 (4.4, 6.2)
<i>p</i> -value <sup>a</sup>	<i>p</i> =0.0002		<i>p</i> =0.0047		<i>p</i> =0.0124		<i>p</i> =0.0002	
Overall Survival (months, 95% CI)	24.8 (22.3,33.7)	20.5 (17.9,25.3)	33.4 (22.8,38.1)	22.8 (18.3,29.8)	22.1 (16.9,33.7)	18.4 (12.7,23.8)	30.5 (26.8, ne)	22.1 (17.6, 28.9)
<i>p</i> -value <sup>a</sup>	<i>p</i> =0.0540		<i>p</i> =0.1021		<i>p</i> =0.2597		<i>p</i> =0.0062	

H = trastuzumab; Chemo = chemotherapy; AC = anthracycline + cyclophosphamide; P = paclitaxel; D = docetaxel  
<sup>a</sup> p = log-rank test; <sup>b</sup> p = Chi-square test, ne = could not be estimated or not yet reached.

### Trastuzumab in Combination with Anastrozole

The TAnDEM trial was a multi-centre, randomized, open-label, phase III trial comparing trastuzumab + anastrozole with anastrozole-alone for the first-line treatment of metastatic breast cancer in HER2 overexpressing, hormone-receptor (i.e. oestrogen-receptor (ER) and/or progesterone-receptor (PR)) positive post-menopausal patients. Two hundred and seven patients were randomized to receive oral anastrozole (1 mg/day) with or without trastuzumab (4 mg/kg loading dose, followed by 2 mg/kg weekly). Patients who had received trastuzumab for early disease were excluded from this trial.

Median progression free survival (PFS) was doubled in the trastuzumab + anastrozole arm compared to the anastrozole-alone arm (4.8 months vs. 2.4 months; *p* = 0.0016). For the other parameters the improvements seen for trastuzumab + anastrozole were; overall response (16.5% vs. 6.7%); clinical benefit rate (42.7% vs. 27.9%); time to progression (4.8 months vs. 2.4 months). For time to response and duration of response no difference could be recorded between the arms. There was no significant difference in overall survival, however more than half of the patients in the anastrozole-alone arm crossed over to a trastuzumab-containing regimen after progression of disease.

### Trastuzumab Monotherapy

Study H0649g was a multinational, multi-centre, single arm trial of trastuzumab as monotherapy in 222 women with HER2 overexpressing metastatic breast cancer. All patients had relapsed following treatment with the best available agents (e.g. anthracyclines and taxanes) and were heavily pre-treated. Two-thirds of the patients had prior adjuvant chemotherapy and all patients had tumour progression following at least one prior regimen of cytotoxic chemotherapy for metastatic disease. Ninety-four percent of the patients had prior anthracycline therapy, approximately 60% had prior paclitaxel therapy and 26% had prior bone marrow or stem cell transplants. Together with HER2 overexpression, which is associated with poorer clinical outcomes, aggressive disease was also suggested by nodal status at diagnosis and by the disease-free interval. Twenty-seven percent of patients had 10 or more positive nodes at the time of diagnosis. Thirty-eight percent of patients had a disease-free interval of less than one year prior to enrolment.

Patients received an intravenous loading dose of 4 mg/kg trastuzumab on Day 0, followed by weekly intravenous infusions of 2 mg/kg until there was evidence of disease progression.

Patients who developed progressive disease could stop treatment, continue on the 2 mg/kg weekly dose or receive an increased intravenous dose of 4 mg/kg, as the investigator deemed appropriate. The primary efficacy parameter was tumour response rate.

Trastuzumab as second- or third-line therapy induced objective, durable tumour responses in women with metastatic breast cancer who had tumours that overexpressed HER2. There were 8 complete responses and 26 partial responses yielding an overall response rate of 15%. The durability of the responses was particularly notable. The median duration of the responses was 9.1 months at the cut-off date for analysis (see Table 11 below).

**Table 11: Efficacy Outcomes with Monotherapy Study H0649g**

Outcome Measure	n	Time (months) Kaplan-Meier Estimate of Median (range)
Duration of response	34	9.1 (2–26+)
Time to disease progression	213	3.1 (0–28+)
Time to Treatment Failure	213	2.4 (0–28+)
Survival Time	213	12.8 (0.5–30+)

The clinical significance of the objective tumour responses in this group of patients was supported by the quality-of-life and survival data. Responders had clinically meaningful improvements in physical function, role function, social function, global quality of life and fatigue scale scores during trastuzumab treatment. Most responders were still alive at data cut-off (28/34; 82%). The Kaplan-Meier estimate of median survival for all treated patients at the data cut-off date was 12.8 months.

Evidence of efficacy for trastuzumab monotherapy is based upon response rates. No data are available to demonstrate improvement in survival or quality of life.

### Advanced Gastric Cancer

Study BO18255 (ToGA) was a randomized, open-label, multicentre phase III trial investigating trastuzumab in combination with a fluoropyrimidine and cisplatin (FP) versus chemotherapy alone as first-line therapy in patients with HER2 positive, inoperable, locally advanced or recurrent and/or metastatic adenocarcinoma of the stomach or gastro-oesophageal junction.

Patients were eligible if they had 3+ levels of HER2 overexpression based on a 0 - 3+ scale by IHC assessment of tumour tissue and/or those whose tumours had HER2 gene amplification as determined

by a FISH test (see DOSAGE AND ADMINISTRATION, Detection of HER2 Overexpression or HER2 Gene Amplification).

After satisfying the screening eligibility criteria, including assessment of HER2 status, patients were randomly assigned (1:1) to receive either trastuzumab (8 mg/kg loading dose, followed by 6 mg/kg every 3 weeks) + fluoropyrimidine/cisplatin (FP+H) or FP alone. The chemotherapy regimen was chosen between 5-FU/cisplatin and capecitabine/cisplatin at the investigator's discretion and could be determined on an individual patient basis.

The efficacy results from ToGA are summarized in Table 12. The primary endpoint was overall survival, defined as the time from the date of randomization to the date of death from any cause. At the time of analysis a total of 349 randomized patients had died: 182 patients (62.8%) in the control arm and 167 patients (56.8%) in the treatment arm. The majority of the deaths were due to events related to the underlying cancer.

Overall survival was significantly improved in the FP + H arm compared to the FP arm ( $p = 0.0046$ , log-rank test). The median survival time was 11.1 months with FP and 13.8 months with FP + H. The risk of death was decreased by 26% (HR = 0.74; 95% CI 0.60 - 0.91) for patients in the FP + H arm compared to the FP arm.

Post-hoc subgroup analyses indicate that targeting tumours with higher levels of HER2 protein (IHC 2+/FISH+ and IHC 3+/regardless of FISH status) results in a greater treatment effect. The median overall survival for the high HER2 expressing group was 11.8 months versus 16 months, HR = 0.65 (95%CI 0.51 - 0.83) and the median PFS was 5.5 months vs. 7.6 months, HR = 0.64 (95% CI 0.51 - 0.79).

**Table 12: Summary of Efficacy from Study BO18255**

Trastuzumab dosage regimen	Every 3 weeks			
<b>Chemotherapy regimens (FP)</b>	<p>Capecitabine: 1000 mg/m<sup>2</sup> orally twice daily for 14 days every 3 weeks for 6 cycles (Days 1 to 15 of each cycle).</p> <p>5-FU: 800 mg/m<sup>2</sup>/day as a continuous IV infusion over 5 days, given every 3 weeks for 6 cycles (Days 1 to 5 of each cycle). The 5-FU infusion could be started at the same time as the cisplatin infusion on Day 1.</p> <p>Cisplatin: 80 mg/m<sup>2</sup> every 3 weeks for 6 cycles (on Day 1 of each cycle) as a 2h IV infusion with hydration and premedication (steroids and anti-emetics).</p>			
<b>Efficacy Parameters</b>	<b>FP = 290</b>	<b>FP+H = 294</b>	<b>HR (95% CI)</b>	<b>-value</b>
Overall Survival, Median months	11.1	13.8	0.74 (0.60-0.91)	0.0046
Progression-Free Survival, Median months	5.5	6.7	0.71 (0.59-0.85)	0.0002
Time to Disease Progression, Median months	5.6	7.1	0.70 (0.58-0.85)	0.0003
Overall Response Rate, %	34.5	47.3	1.70 <sup>a</sup> (1.22, 2.38)	0.0017

Trastuzumab dosage regimen	Every 3 weeks			
Duration of Response, Median months	4.8	6.9	0.54 (0.40-0.73)	<0.0001

FP: fluoropyrimidine/cisplatin; FP+H: fluoropyrimidine/cisplatin + trastuzumab; a Odds ratio

*Progression-free-survival*: time between day of randomization and first documentation of progressive disease (PD) or date of death, whichever occurred first. *Time to disease progression*: time between randomization and first occurrence of PD. *Overall response*: occurrence of either a confirmed complete (CR) or a partial (PR) best overall response as determined by RECIST criteria from confirmed radiographic evaluations of target and non-target lesions. *Duration of response*: time from when response (CR or PR) was first documented to the first documented disease progression. This was only calculated for patients who had a best overall response of CR or PR.

## INDICATIONS

### Early Breast Cancer

OGIVRI is indicated for the treatment of HER2-positive early breast cancer following surgery, and in association with chemotherapy and, if applicable, radiotherapy.

### Locally Advanced Breast Cancer

OGIVRI is indicated for the treatment of HER2-positive locally advanced breast cancer in combination with neoadjuvant chemotherapy followed by adjuvant OGIVRI.

### Metastatic Breast Cancer

OGIVRI is indicated for the treatment of patients with metastatic breast cancer who have tumours that overexpress HER2:

- as monotherapy for the treatment of those patients who have received one or more chemotherapy regimens for their metastatic disease;
- in combination with taxanes for the treatment of those patients who have not received chemotherapy for their metastatic disease; or
- in combination with an aromatase inhibitor for the treatment of post-menopausal patients with hormone-receptor positive metastatic breast cancer.

### Advanced Gastric Cancer

OGIVRI is indicated in combination with cisplatin and either capecitabine or 5-FU for the treatment of patients with HER2 positive advanced adenocarcinoma of the stomach or gastro-oesophageal junction who have not received prior anti-cancer treatment for their metastatic disease.

## CONTRAINDICATIONS

OGIVRI is contraindicated in patients with known hypersensitivity to trastuzumab, Chinese hamster ovary cell proteins or to any of its excipients.



In the treatment of early or locally advanced breast cancer, OGIVRI is contraindicated in patients with a left ventricular ejection fraction of less than 45% and those with symptomatic heart failure.

## PRECAUTIONS

### General

OGIVRI therapy should only be initiated under the supervision of a physician experienced in the treatment of cancer patients. Usual clinical care should be taken to prevent microbial contamination of the intravenous access sites used to deliver OGIVRI therapy. OGIVRI should be administered by a healthcare professional prepared to manage anaphylaxis and adequate life support facilities should be available. Treatment may be administered in an outpatient setting.

If OGIVRI is used concurrently with cytotoxic chemotherapy, the specific guidelines used to reduce or hold the dose of chemotherapy should be followed. Patients may continue OGIVRI therapy during periods of reversible chemotherapy-induced myelosuppression, renal toxicity or hepatic toxicity.

Severe dyspnoea at rest due to complications of advanced malignancy or requiring supplementary oxygen therapy.

### Sorbitol

OGIVRI contains 0.77 mg sorbitol/ 1.00 mg trastuzumab. There is 115.2 mg of sorbitol in the 150 mg vial and 337.9 mg of sorbitol in the 440 mg Pharmacy Bulk Pack.

Patients with hereditary fructose intolerance (HFI) must not be given this medicine.

A detailed history with regards to HFI symptoms has to be taken of each patient prior to being given this medicinal product.

### Cardiac Dysfunction

#### General considerations

Patients treated with trastuzumab are at increased risk of developing congestive heart failure (CHF) (New York Heart Association [NYHA] class II-IV) or asymptomatic cardiac dysfunction. These events have been observed in patients receiving trastuzumab therapy alone or in combination with a taxane following anthracycline (doxorubicin or epirubicin)–containing chemotherapy. This may be moderate to severe and has been associated with death. In addition, caution should be exercised in treating patients with increased cardiac risk e.g. hypertension, documented coronary artery disease, CHF, diastolic dysfunction, older age.

Population pharmacokinetic model simulations indicate that trastuzumab may persist in the circulation for up to 7 months after stopping trastuzumab treatment (see Pharmacokinetics). Patients who receive anthracycline after stopping trastuzumab may also be at increased risk of cardiac dysfunction. If possible, physicians should avoid anthracycline-based therapy for up to 7 months after stopping trastuzumab. If anthracyclines are used, the patient's cardiac function should be monitored carefully.

Candidates for treatment with trastuzumab, especially those with prior anthracycline and cyclophosphamide (AC) exposure, should undergo baseline cardiac assessment including history and physical examination, ECG echocardiogram, and/or MUGA scan. Monitoring may help to identify patients who develop cardiac dysfunction, including signs and symptoms of CHF. Cardiac assessments,

as performed at baseline, should be repeated every 3 months during treatment and every 6 months following discontinuation of treatment until 24 months from the last administration of trastuzumab.

If left ventricular ejection fraction (LVEF) drops 10 percentage points from baseline and to below 50%, trastuzumab should be withheld and a repeat LVEF assessment performed within approximately 3 weeks. If LVEF has not improved, or declined further, or clinically significant CHF has developed, discontinuation of trastuzumab should be strongly considered, unless the benefits for the individual patient are deemed to outweigh the risks.

Patients who develop asymptomatic cardiac dysfunction may benefit from more frequent monitoring (e.g. every 6 - 8 weeks). If patients have a continued decrease in left ventricular function, but remain asymptomatic, the physician should consider discontinuing therapy if no clinical benefit of trastuzumab therapy has been seen.

The safety of continuation or resumption of trastuzumab in patients who experience cardiac dysfunction has not been prospectively studied. If symptomatic cardiac failure develops during trastuzumab therapy, it should be treated with the standard medications for this purpose. In the pivotal trials, most patients who developed heart failure or asymptomatic cardiac dysfunction improved with standard heart failure treatment consisting of angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker -blocker. The majority of patients with cardiac symptoms and evidence of a clinical benefit of trastuzumab treatment continued on weekly therapy with trastuzumab without additional clinical cardiac events.

### **Early and Locally Advanced Breast Cancer**

For patients with early breast cancer, cardiac assessments, as performed at baseline, should be repeated every 3 months during treatment and every 6 months following discontinuation of treatment until 24 months from the last administration of trastuzumab. In patients who receive anthracycline containing chemotherapy further monitoring is recommended and should occur yearly up to 5 years from the last administration of trastuzumab, or longer if a continuous decrease of LVEF is observed.

All patients should have a determination of LVEF prior to treatment. Use of trastuzumab is contraindicated in patients with early or locally advanced disease and a LVEF of less than 45% and those with symptomatic heart failure (see CONTRAINDICATIONS). Patients with a LVEF of 45 - 55% at baseline should be monitored regularly for symptoms of heart failure during trastuzumab treatment.

Patients with history of myocardial infarction (MI), angina pectoris requiring medication, history of or present CHF (NYHA Class II–IV), other cardiomyopathy, cardiac arrhythmia requiring medication, clinically significant cardiac valvular disease, poorly controlled hypertension (hypertension controlled by standard medication eligible), and haemodynamic effective pericardial effusion were excluded from adjuvant and neoadjuvant breast cancer clinical trials with trastuzumab.

### **Adjuvant treatment**

Trastuzumab and anthracyclines should not be given concurrently in the adjuvant treatment setting.

An increase in the incidence of symptomatic and asymptomatic cardiac events was observed when trastuzumab was administered after anthracycline-containing chemotherapy compared to administration with a non-anthracycline regimen of docetaxel and carboplatin. The incidence was more marked when trastuzumab was administered concurrently with taxanes than when administered

sequentially to taxanes. Regardless of the regimen used, most symptomatic cardiac events occurred within the first 18 months.

Risk factors for a cardiac event, identified in 4 large adjuvant studies, included advanced age (> 50 years), low level of baseline and declining LVEF (< 55%), low LVEF prior to or following the initiation of paclitaxel treatment, trastuzumab treatment, and prior or concurrent use of anti-hypertensive medications. In patients receiving trastuzumab after completion of adjuvant chemotherapy the risk of cardiac dysfunction was associated with a higher cumulative dose of anthracycline given prior to initiation of trastuzumab and a high body mass index (> 25 kg/m<sup>2</sup>).

### **Neoadjuvant-adjuvant treatment**

Trastuzumab neoadjuvant-adjuvant treatment concurrent with anthracyclines should be used with caution and only in chemotherapy-naïve patients. The maximum cumulative doses of the low-dose anthracycline regimens should not exceed 180 mg/m<sup>2</sup> (doxorubicin) or 360 mg/m<sup>2</sup> (epirubicin).

If patients have been treated concurrently with low-dose anthracyclines and trastuzumab in the neoadjuvant setting, no additional cytotoxic chemotherapy should be given after surgery.

### **Metastatic breast cancer**

Trastuzumab and anthracyclines should not be given concurrently in the metastatic breast cancer setting.

### **Advanced Gastric Cancer**

In advanced gastric cancer, patients with a history of documented congestive heart failure, angina pectoris requiring medication, evidence of transmural myocardial infarction on ECG, poorly controlled hypertension (systolic BP >180 mmHg or diastolic BP >100 mmHg), clinically significant valvular heart disease, high risk uncontrollable arrhythmias, and baseline LVEF <50% (measured by echocardiography or MUGA) were excluded from Study BO18255 (ToGA) according to the study protocol.

### **Hypersensitivity Reactions including Anaphylaxis**

Severe hypersensitivity reactions have been infrequently reported in patients treated with trastuzumab. Signs and symptoms include anaphylaxis, urticaria, bronchospasm, angioedema, and/or hypotension. In some cases, the reactions have been fatal. The onset of symptoms generally occurred during an infusion, but there have also been reports of symptom onset after the completion of an infusion. Reactions were most commonly reported in association with the initial infusion.

Patients should be observed closely for hypersensitivity reactions, OGIVRI infusion should be interrupted in all patients with severe hypersensitivity reactions. In the event of a hypersensitivity reaction, appropriate medical therapy should be administered, which may include adrenaline, corticosteroids, antihistamines, bronchodilators and oxygen. Patients should be evaluated and carefully monitored until complete resolution of signs and symptoms.

On very rare occasions, patients have experienced the onset of infusion symptoms and pulmonary symptoms more than six hours after the start of the trastuzumab infusion. Patients should be warned of the possibility of such a late onset and should be instructed to contact their physician if these symptoms occur.

## Infusion-Related Reactions (IRRs)

IRRs are known to occur with the administration of trastuzumab (See ADVERSE EFFECTS).

Pre-medication may be used to reduce risk of occurrence of IRRs.

Serious IRRs to trastuzumab infusion including dyspnoea, hypotension, wheezing, bronchospasm, tachycardia, reduced oxygen saturation and respiratory distress and supraventricular tachyarrhythmia have been reported (see ADVERSE EFFECTS).

Patients should be observed for IRRs. Interruption of an IV infusion may help control such symptoms and the infusion may be resumed when symptoms abate. These symptoms can be treated with an analgesic/antipyretic such as paracetamol and an antihistamine. Serious reactions have been treated successfully with supportive therapy such as oxygen, intravenous fluids, beta-agonists and corticosteroids. In rare cases, these reactions are associated with a clinical course culminating in a fatal outcome. In other patients with acute onset of signs and symptoms, initial improvement was followed by clinical deterioration and delayed reactions with rapid clinical deterioration have also been reported. Fatalities have occurred within hours or up to one week following an infusion.

Patients who are experiencing dyspnoea at rest due to complications of advanced malignancy or co-morbidities may be at increased risk of a fatal infusion reaction. Therefore, these patients should not be treated with trastuzumab (see Pulmonary Reactions).

## Pulmonary Reactions

Severe pulmonary events leading to death have been reported with the use of trastuzumab in the post-marketing setting. These events may occur as part of an infusion-related reaction (see Infusion-Related Reactions) or with a delayed onset. In addition, cases of interstitial lung disease including lung infiltrates, acute respiratory distress syndrome, pneumonia, pneumonitis, pleural effusion, respiratory distress, acute pulmonary oedema, pulmonary hypertension, pulmonary fibrosis and respiratory insufficiency have been reported.

Risk factors associated with interstitial lung disease include prior or concomitant therapy with other anti-neoplastic therapies known to be associated with it such as taxanes, gemcitabine, vinorelbine and radiation therapy. Patients with dyspnoea at rest due to complications of advanced malignancy and co-morbidities may be at increased risk of pulmonary events. Therefore, these patients should not be treated with trastuzumab.

## Ability to Drive and Use Machines

No studies on the effects on the ability to drive and to use machines have been performed. Patients experiencing infusion-related symptoms should be advised not to drive or use machines until symptoms resolve completely.

## Effects on Fertility

A study in female cynomolgus monkeys revealed no evidence of impaired fertility at IV trastuzumab doses up to 25 mg/kg twice weekly, corresponding to serum trough levels (serum  $C_{min}$ ) about 15 times higher than that in humans receiving the recommended weekly dose of 2 mg/kg.

## Use in Pregnancy – Category D

OGIVRI should be avoided during pregnancy and since trastuzumab may persist in the circulation for up to 7 months, pregnancy should be avoided for 7 months after the last dose of OGIVRI, unless the anticipated benefit for the mother outweighs the unknown risk to the foetus.

In the post-marketing setting, cases of foetal renal growth and/or function impairment in association with oligohydramnios, some associated with fatal pulmonary hypoplasia of the foetus, have been reported in pregnant women receiving OGIVRI.

Women of childbearing potential should be advised to use effective contraception during treatment with OGIVRI and for at least 7 months after treatment has concluded. Women who become pregnant should be advised of the possibility of harm to the foetus. If a pregnant woman is treated with OGIVRI, or becomes pregnant within 7 months following the last dose of OGIVRI, close monitoring by a multidisciplinary team is desirable.

## Use in Lactation

A study conducted in lactating cynomolgus monkeys dosed intravenously with trastuzumab at 25 mg/kg twice weekly (serum  $C_{min}$  about 15 times higher than that in humans receiving the recommended weekly dose of 2 mg/kg) demonstrated that trastuzumab is excreted in the milk. The presence of trastuzumab in the serum of infant monkeys was not associated with adverse effects on their growth or development from birth to 1 month of age. However, the binding affinity of trastuzumab to epidermal growth factor receptor 2 protein in cynomolgus monkeys is unclear.

It is not known whether trastuzumab is excreted in human milk. As human immunoglobulin G (IgG) is secreted into human milk and the potential for harm to the infant is unknown, breast-feeding should be avoided during trastuzumab therapy and for 7 months after the last dose of trastuzumab.

## Paediatric Use

The safety and efficacy of trastuzumab in patients under the age of 18 years have not been established.

## Use in the Elderly

Clinical experience is limited in patients above 65 years of age. The risk of cardiac dysfunction may be increased in elderly patients. The reported clinical experience is not adequate to determine whether older patients respond differently from younger patients. Elderly patients did not receive reduced doses of trastuzumab in clinical trials. However, greater sensitivity to trastuzumab in some older patients cannot be ruled out.

## Genotoxicity

Trastuzumab did not induce gene mutations in bacteria, nor did it cause chromosomal damage *in vitro* (chromosome aberration assay in human lymphocytes) or *in vivo* (mouse micronucleus test).

## Carcinogenicity

No studies on the carcinogenic potential of trastuzumab have been conducted to date.

## Use in Renal Impairment

Formal PK studies have not been conducted in patients with renal impairment. Based on population PK analysis, renal impairment is not expected to influence trastuzumab exposure, however, limited data from patients with moderate to severe renal impairment were included in the population PK analysis (see Pharmacokinetics).

## Use in Hepatic Impairment

The use of trastuzumab in patients with hepatic impairment has not been studied.

## INTERACTIONS WITH OTHER MEDICINES

No formal drug interaction studies have been performed with trastuzumab in humans. Clinically significant interactions with concomitant medication used in clinical trials have not been observed. A comparison of serum levels of trastuzumab given in combination with cisplatin, doxorubicin or epirubicin-plus-cyclophosphamide has not suggested the possibility of any interaction.

Administration of paclitaxel in combination with trastuzumab resulted in a slightly less than two-fold decrease in trastuzumab clearance in a non-human primate study and a 1.5-fold increase in trastuzumab serum levels in clinical studies. Paclitaxel pharmacokinetics determined during the fourth cycle of the alternative 3-weekly trastuzumab regimen (n = 25) were not altered appreciably, relative to parameters determined during the initiation of paclitaxel, prior to introduction of trastuzumab. Similarly, docetaxel pharmacokinetics determined during the first dose of trastuzumab in the standard weekly regimen (n = 10) were not altered appreciably relative to those determined 2 weeks earlier for docetaxel-alone.

A pharmacokinetic interaction substudy of BO18255 (ToGA) performed in male and female Japanese patients with advanced gastric cancer showed that co-administration of trastuzumab and capecitabine and cisplatin had no significant effects on the pharmacokinetics of the two chemotherapy agents compared with co-administration of the two agents without trastuzumab. The pharmacokinetics of trastuzumab were not evaluated in this study.

The administration of concomitant chemotherapy (either anthracycline or cyclophosphamide or anastrozole) did not appear to influence the pharmacokinetics of trastuzumab.

## ADVERSE EFFECTS

The adverse drug reactions listed in this section fall into the following categories: very common ( 1/10); common ( <1/10,000); not known (cannot be estimated from the available data).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Presented in Table 13 below are adverse reactions that have been reported in association with the use of trastuzumab alone, or in combination with chemotherapy in the below pivotal clinical trials as well as in the post-marketing setting.

### Early Breast Cancer

BO16348 (HERA): trastuzumab arm (n=1678). Control arm (n=1708)



B-31/N9831 Joint Analysis: trastuzumab arms (n=2345). Control arm (n=1673)

BCIRG 006: trastuzumab arm (n=2133). Control arm (n=1041)

BO16216 (TanDEM): trastuzumab arm (n=161). Control arm (n=161)

### Locally Advanced Breast Cancer

MO16432 (NOAH): trastuzumab arm (n=115). Control arm (n=116)

### Metastatic Breast Cancer (MBC)

H0648g / H0649g: trastuzumab arms ( n=469 and n=222 respectively)

M77001: trastuzumab arm (n=92). Control arm (n=94).

### Advanced Gastric Cancer

BO18255 (ToGA): trastuzumab arm (n=294). Control arm (n=290)

All terms included are based on the highest percentage seen in pivotal clinical trials.

**Table 13: Adverse Reactions**

System organ class	Adverse reaction <sup>1</sup>	Frequency
Infections and infestations	Nasopharyngitis	Very common
	Infection	Very common
	Neutropenic sepsis	Common
	Cystitis	Common
	Herpes zoster	Common
	Influenza	Common
	Pharyngitis	Common
	Sinusitis	Common
	Skin infection	Common
	Rhinitis	Common
	Upper respiratory tract infection	Common
	Urinary tract infection	Common
	Erysipelas	Common
	Cellulitis	Common
	Sepsis	Uncommon
Neoplasms benign, malignant and unspecified (incl. Cysts and polyps)	Malignant neoplasm progression	Not known
	Neoplasm progression	Not known
Blood and lymphatic system disorders	Febrile neutropenia	Very common
	Anaemia	Very common
	Thrombocytopenia	Very common
	White blood cell count decreased / leukopenia	Very common
	Neutropenia	Very common
	Hypoprothrombinaemia	Not known
	Immune Thrombocytopenia	Not Known
Immune system disorders	Hypersensitivity	Common
	<sup>2</sup> Anaphylactic reaction	Not known
	<sup>2</sup> Anaphylactic shock	Not known
Metabolism and nutrition disorders	Weight Decreased/Weight Loss	Very common
	Weight Increased	Very common
	Decreased appetite	Very common

OGIVRI – PRODUCT INFORMATION

22

System organ class	Adverse reaction <sup>1</sup>	Frequency
Psychiatric disorders	Anorexia	Very common
	Hyperkalaemia	Not known
	Insomnia	Very common
	Depression	Common
	Anxiety	Common
Nervous system disorders	Thinking abnormal	Common
	Tremor <sup>3</sup>	Very common
	Dizziness	Very common
	Headache	Very common
	Dysgeusia	Very common
	Paraesthesia	Very common
	Hypoaesthesia	Very common
	Peripheral neuropathy	Common
	Hypertonia	Common
	Somnolence	Common
	Ataxia	Common
	Paresis	Rare
	Brain oedema	Not known
Eye disorders	Conjunctivitis	Very common
	Lacrimation increased	Very common
	Dry eye	Common
	Papilloedema	Not known
	Retinal haemorrhage	Not known
Ear and Labyrinth Disorders	Deafness	Uncommon
Cardiac disorders	<sup>3</sup> Blood pressure decreased	Very common
	<sup>3</sup> Blood pressure increased	Very common
	<sup>3</sup> Heart beat irregular	Very common
	<sup>3</sup> Palpitation	Very common
	<sup>3</sup> Cardiac flutter	Very common
	<sup>4</sup> Ejection fraction decreased	Very common
	<sup>2</sup> Cardiac failure (congestive)	Common
	<sup>2,3</sup> Supraventricular tachyarrhythmia	Common
	Cardiomyopathy	Common
	Pericardial effusion	Uncommon
	Cardiogenic shock	Not known
	Pericarditis	Not known
	Bradycardia	Not known
	Gallop rhythm present	Not known
Vascular disorders	Lymphoedema	Very common
	Hot flush	Very common
	<sup>2,3</sup> Hypotension	Common
	Hypertension	Common
	Vasodilatation	Common
Respiratory, thoracic and mediastinal disorders	<sup>2,3</sup> Wheezing	Very common
	<sup>2</sup> Dyspnoea	Very common
	Cough	Very common
	Epistaxis	Very common
	Rhinorrhoea	Very common
	Oropharyngeal pain	Very common
	Asthma	Common
	Lung disorder	Common
	<sup>2</sup> Pleural effusion	Common
	<sup>2</sup> Pneumonia	Common
	Pneumonitis	Uncommon

System organ class	Adverse reaction <sup>1</sup>	Frequency
	<sup>2</sup> Pulmonary fibrosis	Not known
	<sup>2</sup> Respiratory distress	Not known
	<sup>2</sup> Respiratory failure	Not known
	<sup>2</sup> Lung infiltration	Not known
	<sup>2</sup> Acute pulmonary oedema	Not known
	<sup>2</sup> Acute respiratory distress syndrome	Not known
	<sup>2</sup> Bronchospasm	Not known
	<sup>2</sup> Hypoxia	Not known
	<sup>2</sup> Oxygen saturation decreased	Not known
	Laryngeal oedema	Not known
	<sup>2</sup> Orthopnoea	Not known
	Pulmonary oedema	Not known
	Interstitial lung disease	Not known
Gastrointestinal disorders	Diarrhoea	Very common
	Vomiting	Very common
	Nausea	Very common
	Lip swelling	Very common
	Abdominal pain	Very common
	Stomatitis	Very common
	Pancreatitis	Very common
	Constipation	Very common
	Dyspepsia	Very common
	Haemorrhoids	Common
	Dry mouth	Common
Hepatobiliary disorders	Hepatocellular Injury	Common
	Hepatitis	Common
	Liver Tenderness	Common
	Jaundice	Rare
	Hepatic Failure	Not known
Skin and subcutaneous disorders	Erythema	Very common
	Rash	Very common
	Swelling face <sup>3</sup>	Very common
	Palmar-plantar erythrodysaesthesia syndrome	Very common
	Nail disorder	Very common
	Alopecia	Very common
	Dry skin	Common
	Ecchymosis	Common
	Hyperhydrosis	Common
	Maculopapular rash	Common
	Acne	Common
	Onychoclasia	Common
	Pruritus	Common
	Dermatitis	Common
	Urticaria	Uncommon
	Angioedema	Not known
Musculoskeletal and connective tissue disorders	Arthralgia	Very common
	Muscle tightness	Very common
	Myalgia	Very common
	Arthritis	Common
	Back pain	Common
	Bone pain	Common
	Muscle spasms	Common

System organ class	Adverse reaction <sup>1</sup>	Frequency
Renal and urinary conditions	Neck pain	Common
	Pain in extremity	Common
	Renal disorder	Common
	Glomerulonephritis membranous	Not known
	Glomerulonephropathy	Not known
Pregnancy, puerperium and perinatal disorders	Renal failure	Not known
	Oligohydramnios	Not known
	Renal hypoplasia	Not known
	Pulmonary hypoplasia	Not known
Reproductive system and breast disorders	Breast inflammation/mastitis	Common
General disorders and administration site conditions	Asthenia	Very common
	Chest pain	Very common
	Chills	Very common
	Fatigue	Very common
	Influenza-like symptoms	Very common
	Infusion related reaction	Very common
	Pain	Very common
	Pyrexia	Very common
	Peripheral oedema	Very common
	Mucosal inflammation	Very common
	Malaise	Common
	Oedema	Common
Injury, poisoning and procedural complications	Nail toxicity	Very common
	Contusion	Common

<sup>1</sup> Adverse drug reactions (ADRs) were identified as events that occurred with at least a 2% difference compared to the control arm in at least one of the major randomised clinical trials; <sup>2</sup> Denotes adverse reactions that have been reported in association with a fatal outcome; <sup>3</sup> Denotes adverse reactions that are reported largely in association with Infusion-related reactions. Specific percentages for these are not available; <sup>4</sup> Observed with combination therapy following anthracyclines and combined with taxanes

The following information is relevant to all indications.

### Infusion-Related Reactions (IRRs) and Hypersensitivity

IRRs such as chills and/or fever, dyspnoea, hypotension, wheezing, bronchospasm, tachycardia, reduced oxygen saturation and respiratory distress were seen in all trastuzumab clinical trials (see PRECAUTIONS).

IRRS may be clinically difficult to distinguish from hypersensitivity reactions.

The rate of IRRs of all grades varied between studies depending on the indication, whether trastuzumab was given concurrently with chemotherapy or as monotherapy and data collection methodology.

In early breast cancer, the rate of IRRs ranged from 18% to 54% in the trastuzumab containing arm compared to 6% to 50% in the comparator arm (which may have contained other chemotherapy). Severe reactions (grade 3 and above) ranged from 0.5% to 6% in the trastuzumab containing arm compared to 0.3% to 5 % in the comparator arm.

In metastatic breast cancer, the rate of IRRs ranged from 49% to 54% in the trastuzumab containing arm compared to 36% to 58% in the comparator arm (which may have contained other chemotherapy),

Severe reactions (grade 3 and above) ranged from 5 % to 7% in the trastuzumab containing arm compared to 5% to 6% in the comparator arm.

Anaphylactoid reactions were observed in isolated cases (see PRECAUTIONS).

### **Cardiac Dysfunction**

Congestive heart failure (NYHA Class II-IV) is a common adverse reaction to trastuzumab. It has been associated with fatal outcome. Signs and symptoms of heart failure, such as dyspnoea, orthopnoea, increased cough, pulmonary oedema, pulmonary hypertension and S3 gallop or reduced ventricular ejection fraction, have been observed in patients treated with trastuzumab (see PRECAUTIONS).

### **Locally Advanced Breast Cancer (neoadjuvant –adjuvant setting)**

In the clinical trial setting, when trastuzumab was administered concurrently with neoadjuvant chemotherapy containing 3-4 cycles of a neoadjuvant anthracycline (cumulative doxorubicin dose 180 mg/m<sup>2</sup> or epirubicin dose 360 mg/m<sup>2</sup>) overall, the incidence of symptomatic cardiac dysfunction was up to 1.7 % in the trastuzumab arm.

### **Early Breast Cancer (adjuvant setting)**

In 3 pivotal clinical trials of adjuvant trastuzumab given in combination with chemotherapy the incidence of grade 3/4 cardiac dysfunction (symptomatic CHF) was similar in patients who were administered chemotherapy alone and in patients who were administered trastuzumab sequentially to a taxane (0.3 - 0.4%). The rate was highest in patients who were administered trastuzumab concurrently

cyclophosphamide followed by paclitaxel) + H (trastuzumab) was estimated at 3.2%, compared with n the cumulative incidence of cardiac events was seen with further follow-up at 5 years.

At 5.5 years, the rates of symptomatic cardiac or LVEF events were 1.0%, 2.3%, and 1.1% in the (doxorubicin plus cyclophosphamide, followed by docetaxel plus trastuzumab), and DCarbH (docetaxel, carboplatin and trastuzumab) treatment arms, respectively. For symptomatic CHF (NCI-CTC Grade 3 - 4), the 5-year

developing a symp  
continuous increase in the cumulative rate of symptomatic cardiac or LVEF events up to 2.3% compared

When trastuzumab was administered after completion of adjuvant chemotherapy, NYHA class III-IV heart failure was observed in 0.6% of patients in the 1 year arm after a median follow up of 12 months. After a median follow-up of 3.6 years the incidence of severe CHF and left ventricular dysfunction after 1 year trastuzumab therapy remained low at 0.8% and 9.8%, respectively.

After a median follow-up of 8 years the incidence of severe CHF (NYHA Class III & IV) following 1 year of trastuzumab therapy (combined analysis of the two trastuzumab treatment arms) was 0.8%, and the rate of mild symptomatic and asymptomatic left ventricular dysfunction was 4.6%.

Reversibility of severe CHF (defined as a sequence of at least two consecutive LVEF values 50% after the event) was evident for 71.4% of trastuzumab-treated patients. Reversibility of mild symptomatic and asymptomatic left ventricular dysfunction was demonstrated for 79.5% of

trastuzumab-treated patients. Approximately 17% of cardiac dysfunction related events occurred after completion of trastuzumab.

In the joint analysis of studies NSABP B-31 and NCCTG N9831, with a median follow-up of 8.1 years

in patients with a symptomatic CHF event, while data are missing for 22.6%, 64.5% were known to recover, and 12.9% experienced no recovery. The median time to first recovery by LVEF status occurred at 8.3 months (range 1 – 104 months); 90.3% experienced a full or partial LVEF recovery.

### Metastatic Breast Cancer

Depending on the criteria used to define cardiac dysfunction, the incidence in the pivotal metastatic trials varied between 9% and 12% in the trastuzumab + paclitaxel subgroup, compared with 1% - 4% for the paclitaxel-alone subgroup. For trastuzumab monotherapy, the rate was 6 - 9%. The highest rate of cardiac dysfunction was seen in patients receiving concurrent trastuzumab + anthracycline / cyclophosphamide (27%), significantly higher than in the anthracycline / cyclophosphamide-alone subgroup (7 - 10%). In study M77001 with prospective monitoring of cardiac function, the incidence of symptomatic heart failure was 2.2% in patients receiving trastuzumab and docetaxel, compared with 0% in patients receiving docetaxel-alone. Most of the patients (79%) who developed cardiac dysfunction in these trials experienced an improvement after receiving standard treatment for heart failure.

### Advanced Gastric Cancer

In Study BO18255 (ToGA), at screening, the median LVEF value was 64% (range 48% - 90%) in the fluoropyrimidine/cisplatin (FP) arm and 65% (range 50% - 86%) in the trastuzumab + FP arm.

The majority of the LVEF decreases noted in Study BO18255 (ToGA) were asymptomatic, with the exception of 1 patient in the trastuzumab arm whose LVEF decrease coincided with cardiac failure.

**Table 14: Summary of LVEF Change from baseline (Study BO18255)**

<b>LVEF Decrease<sup>#</sup>: Lowest Post-screening Value</b>	<b>FP ( = 290) (% patients in each treatment arm)</b>	<b>FP + H ( = 294) (% patients in each treatment arm)</b>
10% to <50%	1.1%	4.6%
Absolute Value <50%	1.1%	5.9%
50%	11.8%	16.5%

FP: fluoropyrimidine/cisplatin; FP+H: fluoropyrimidine/cisplatin + trastuzumab; <sup>#</sup>Only includes patients whose method of assessment at that visit is the same as at their initial assessments (FP: n = 187 and FP + H: n = 237).

	<b>FP ( = 290) (% patients in each treatment arm)</b>	<b>FP + H ( = 294) (% patients in each treatment arm)</b>
Total Cardiac Events	6%	6%
Grade 3 NCI CTCAE v3.0	3% <sup>a</sup>	1% <sup>b</sup>

FP: fluoropyrimidine/cisplatin; FP+H: fluoropyrimidine/cisplatin + trastuzumab; <sup>a</sup> 9 patients experienced 9 Events; <sup>b</sup> 4 patients experienced 5 Events

Overall, there were no significant differences in cardiotoxicity between the treatment arm and the comparator arm.



## Haematological Toxicity

Febrile neutropenia, leukopenia, anaemia, thrombocytopenia and neutropenia occurred very commonly. The frequency of occurrence of hypoprothrombinemia is not known. The risk of neutropenia may be slightly increased when trastuzumab is administered with docetaxel following anthracycline therapy.

### Breast Cancer

#### Monotherapy– Study H0649g

Haematological toxicity is infrequent following the administration of trastuzumab as monotherapy in the metastatic setting, WHO Grade 3 leucopenia, thrombocytopenia and anaemia occurring in <1% of patients. No WHO Grade 4 toxicities were observed.

#### Combination Therapy – Studies H0648g and M77001

WHO Grade 3 or 4 haematological toxicity was observed in 63% of patients treated with trastuzumab and an anthracycline/cyclophosphamide compared to an incidence of 62% in patients treated with the anthracycline/cyclophosphamide combination without trastuzumab.

There was an increase in WHO Grade 3 or 4 haematological toxicity in patients treated with the combination of trastuzumab and paclitaxel compared with patients receiving paclitaxel-alone (34% vs. 21%). Haematological toxicity was also increased in patients receiving trastuzumab and docetaxel, compared with docetaxel-alone (32% grade 3/4 neutropenia vs. 22%, using NCI-CTC criteria). The incidence of febrile neutropenia/neutropenic sepsis was also increased in patients treated with trastuzumab + docetaxel (23% vs. 17% for patients treated with docetaxel-alone).

#### Early Setting – HERA Trial

Using NCI-CTC criteria, in the BO16348 (HERA) trial, 0.4% of trastuzumab treated patients experienced a shift of 3 or 4 grades from baseline, compared with 0.6% in the observation arm.

### Advanced Gastric Cancer

The most frequently reported adverse events categorized under the Blood and Lymphatic System Disord

**Table 16: Frequently reported adverse events grade > 3 in blood and lymphatic System Disorders (SOC)**

	<b>FP</b> <b>( = 290)</b> <b>(% patients in each treatment</b> <b>arm)</b>	<b>FP + H</b> <b>( = 294)</b> <b>(% patients in each treatment</b> <b>arm)</b>
Neutropenia	30%	27%
Anaemia	10%	12%
Febrile neutropenia	3%	5%
Thrombocytopenia	3%	5%

FP: fluoropyrimidine/cisplatin; FP+H: fluoropyrimidine/cisplatin + trastuzumab

The total percentage of patients who experienced an adverse event of Grade 3 NCI CTCAE v3.0 categorized under this SOC were 38% in the FP arm and 40% in the FP + H arm.

Overall, there were no significant differences in haematotoxicity between the treatment arm and the comparator arm.

## **Hepatic and Renal Toxicity**

### **Breast Cancer**

#### Monotherapy– Study H0649g

WHO Grade 3 or 4 hepatic toxicity was observed in 12% of patients following administration of trastuzumab as monotherapy in the metastatic setting. This toxicity was associated with progression of disease in the liver in 60% of these patients. No WHO Grade 3 or 4 renal toxicity was observed.

#### Combination Therapy – Study H0648g

WHO Grade 3 or 4 hepatic toxicity was observed in 6% of patients treated with trastuzumab and an anthracycline/cyclophosphamide compared with an incidence of 8% in patients treated with the anthracycline/cyclophosphamide combination without trastuzumab. No WHO Grade 3 or 4 renal toxicity was observed.

WHO Grade 3 or 4 hepatic toxicity was less frequently observed among patients receiving trastuzumab and paclitaxel than among patients receiving paclitaxel-alone (7% vs.15%). No WHO Grade 3 or 4 renal toxicity was observed.

### **Advanced Gastric Cancer**

In Study BO18255 (ToGA) no significant differences in hepatic and renal toxicity were observed between the two treatment arms.

#### NCI-CTCAE (v

trastuzumab than those in the fluoropyrimidine/cisplatin arm (3% and 2% respectively).

#### NCI-

was the only reported adverse event and was not significantly higher in patients receiving trastuzumab than those in the fluoropyrimidine/cisplatin arm (1% and <1% respectively).

rubinaemia

## **Diarrhoea**

### **Breast Cancer**

#### Monotherapy– Study H0649g

Of patients treated with trastuzumab monotherapy in the metastatic setting, 27% experienced diarrhoea.

#### Combination Therapy – Studies H0648g and M77001

An increase in the incidence of diarrhoea, primarily mild to moderate in severity, has been observed in patients receiving trastuzumab in combination with chemotherapy compared with patients receiving chemotherapy-alone or trastuzumab-alone.

#### Early Setting – HERA Study

In the HERA trial, 8% of trastuzumab treated patients experienced diarrhoea during the first year of treatment.

### **Advanced Gastric Cancer**

In Study BO18255 (ToGA), 109 patients (37%) in the trastuzumab treatment arm versus 80 patients (28%) in the comparator arm experienced any grade diarrhoea. Four percent (4%) of patients in the fluoropyrimidine/cisplatin arm experienced Grade  $\geq 3$  diarrhoea vs. 9% in the trastuzumab arm.

### **Infection**

An increased incidence of infections, primarily mild upper respiratory infections of minor clinical significance or catheter infections, has been observed primarily in patients treated with trastuzumab + chemotherapy compared with patients receiving chemotherapy-alone or trastuzumab-alone.

### **Laboratory Abnormalities**

Febrile neutropenia occurs very commonly. Commonly occurring adverse reactions include anaemia, leukopenia, thrombocytopenia and neutropenia. The frequency of occurrence of hypoprothrombinemia is not known.

### **Immunogenicity**

In a neoadjuvant-adjuvant breast cancer trial, 8.1% (24/296) of patients treated with trastuzumab developed antibodies against trastuzumab (regardless of antibody presence at baseline). Neutralizing anti-trastuzumab antibodies were detected in post-baseline samples in 2 of 24 trastuzumab patients.

The clinical relevance of these antibodies is not known. However, the pharmacokinetics, efficacy [determined by pathological complete response (pCR)] or safety [determined by the occurrence of administration related reaction (ARRs)] of trastuzumab did not appear to be adversely affected by these antibodies.

### **Switching treatment from Trastuzumab IV to Trastuzumab SC and vice versa**

Study MO22982 (PrefHER) investigated switching from Trastuzumab IV to Trastuzumab SC, and vice versa, in patients with HER2 positive EBC, with a primary objective to evaluate patient preference for either Trastuzumab IV infusion or Trastuzumab SC injection. This trial investigated using a 2-arm, cross-over design with patients being randomized to one of two different q3w Trastuzumab treatment sequences (Trastuzumab IV (Cycles 1- Trastuzumab SC (Cycles 5-8), or Trastuzumab SC (Cycles 1- Trastuzumab IV (Cycles 5-8)). Patients participating in this trial could be enrolled at any time as long as there were at least 10 remaining cycles of Trastuzumab in their planned treatment regimen, therefore patients were either naïve to Trastuzumab IV treatment (20.3%) or pre-exposed to Trastuzumab IV (79.7%) as part of ongoing adjuvant treatment for HER2 positive EBC. Overall, switches from Trastuzumab IV to Trastuzumab SC and vice versa were well tolerated. Pre-switch rates (Cycles 1-4) for SAEs, Grade 3 AEs and treatment discontinuations due to AEs were low (<5%) and similar to post-switch rates (Cycles 5-8). No Grade 4 or Grade 5 AEs were reported. The effect of multiple switches back and forth was not investigated (see also CLINICAL TRIALS).

## **DOSAGE AND ADMINISTRATION**

HER2 testing is mandatory prior to initiation of trastuzumab therapy (refer to Detection of HER2 Protein Overexpression and Gene Amplification below).

In order to prevent medication errors, it is important to check the vial labels to ensure the medicine being prepared and administered is **Trastuzumab** (OGIVRI) and not **Trastuzumab emtansine** (KADCYLA<sup>®</sup>).

It is important to check the labels to ensure the correct formulation (intravenous or subcutaneous) is being administered to the patient as was prescribed. Switching treatment between Trastuzumab IV and Trastuzumab SC and vice versa, using a three-weekly (q3w) dosing regimen, was investigated in study MO22982 (PrefHER) (see CLINICAL TRIALS, ADVERSE EFFECTS).

In order to improve traceability of biological medicinal products, the trade name and the batch number of the administered product should be clearly recorded in the patient medical record.

Do not administer OGIVRI to patients with rare hereditary problems of fructose intolerance because it contains sorbitol as an excipient (see Precautions).

## Recommended Dosage

### Early Breast Cancer

*Three-weekly schedule:* the recommended initial loading dose is 8 mg/kg body weight, followed by a maintenance dose of 6 mg/kg body weight administered at 3 weekly intervals.

*Weekly schedule:* the recommended initial loading dose is 4 mg/kg body weight, followed by a maintenance dose of 2 mg/kg body weight administered at weekly intervals.

### Locally Advanced Breast Cancer

*Three-weekly schedule:* the recommended initial loading dose is 8 mg/kg body weight, followed by a maintenance dose of 6 mg/kg body weight administered at 3 weekly intervals.

### Metastatic Breast Cancer

*Three-weekly schedule:* the recommended initial loading dose is 8 mg/kg body weight, followed by a maintenance dose of 6 mg/kg body weight administered at 3 weekly intervals.

*Weekly schedule:* the recommended initial loading dose is 4 mg/kg body weight, followed by a maintenance dose of 2 mg/kg body weight administered at weekly intervals.

### Advanced Gastric Cancer

*Three-weekly schedule:* the recommended initial loading dose is 8 mg/kg body weight, followed by a maintenance dose of 6 mg/kg body weight administered at 3-weekly intervals.

Refer to the CLINICAL TRIALS section (including Table 2 for early breast cancer) for the sequence and dosing of chemotherapy medicines used in the supporting pivotal trials. Refer also to the currently approved product information for the chemotherapy partners.

## Administration

OGIVRI IV solution is not to be used for subcutaneous administration and must be administered as an IV infusion. Do not administer as an IV push or bolus.

OGIVRI IV loading doses should be administered over approximately 90 minutes. If the loading dose was well tolerated, subsequent doses can be administered as a 30 minute infusion.

Patients should be observed for fever and chills or other infusion-associated symptoms (see ADVERSE EFFECTS). Interruption of the infusion and/or medication may help to control such symptoms. The infusion may be resumed when symptoms abate.

### Duration of Treatment

Patients with **early or locally advanced breast cancer** should be treated for 1 year or until disease recurrence, or unmanageable toxicity, whichever occurs first. However, extending adjuvant treatment beyond one year is not recommended (see CLINICAL TRIALS, Early breast cancer).

Patients with **metastatic breast cancer** and **advanced gastric cancer** should be treated until progression of disease.

### Missed Doses

If the patient has missed a dose of OGIVRI by one week or less, then the usual maintenance dose of OGIVRI (weekly regimen: 2 mg/kg; 3-weekly: 6 mg/kg) should be administered as soon as possible (do not wait until the next planned cycle). Subsequent maintenance doses should then be administered 7 days or 21 days later according to the weekly or three-weekly schedules, respectively.

If the patient has missed a dose of OGIVRI by more than one week, a re-loading dose of OGIVRI should be administered over approximately 90 minutes (weekly regimen: 4 mg/kg; 3-weekly: 8 mg/kg) as soon as possible. Subsequent maintenance doses (weekly regimen: 2 mg/kg; 3-weekly: 6 mg/kg) should be administered 7 days or 21 days later according to the weekly or three-weekly schedules, respectively.

### Dose Reduction

If the patient develops an infusion-related reaction (IRR), the infusion rate of OGIVRI may be slowed or interrupted (see Precautions). No reductions in the dose of trastuzumab were made during clinical trials. Patients may continue trastuzumab therapy during periods of reversible, chemotherapy-induced myelosuppression, but they should be carefully monitored for complications of neutropenia during this time. The specific instructions to reduce or hold the dose of chemotherapy should be followed.

*Use in Elderly:* In clinical trials, elderly patients did not receive reduced doses of trastuzumab. Age has been shown to have no effect on the disposition of trastuzumab (see PHARMACOKINETICS, Pharmacokinetics in Special Populations).

### Detection of HER2 Protein Overexpression or HER2 Gene Amplification

OGIVRI should only be used in patients whose tumours have HER2 protein overexpression or HER2 gene amplification.

To ensure accurate and reproducible results, testing must be performed in a specialized laboratory, which can ensure validation of the testing procedures.

HER2 protein overexpression should be detected using an immunohistochemistry (IHC)-based assessment of fixed tumour blocks. HER2 gene amplification should be detected using in situ hybridization (ISH) of fixed tumour blocks. Examples of ISH include fluorescence in situ hybridization (FISH), chromogenic in situ hybridization (CISH) and silver in situ hybridization (SISH).

For any other method to be used for the assessment of HER2 protein or gene expression, the test method must be precise and accurate enough to demonstrate overexpression of HER2 (it must be able to distinguish between moderate (congruent with 2+) and strong (congruent with 3+) HER2 overexpression).

For full instructions on the use of these assays and interpretation of the results please refer to the package inserts of validated FISH, CISH and SISH assays. Official recommendations on HER2 testing may also apply.

### **Breast Cancer**

OGIVRI treatment is only appropriate if there is strong HER2 overexpression, as described by a 3+ score by IHC or a positive ISH result. For patients with an intensity score of 2+ on IHC, confirmation of HER2 positive status by ISH is mandatory.

### **Advanced Gastric Cancer**

OGIVRI treatment is only appropriate if there is HER2 overexpression, as described by a 3+ IHC score. For cases with a score of less than 3+ by IHC, confirmation of HER2 positive status by ISH is mandatory.

Bright-field ISH technology is recommended for advanced gastric cancer samples to enable evaluation of tumour histology and morphology in parallel. Either FISH or SISH are recommended for detecting HER2 gene amplification in advanced gastric cancer tissue.

### **Preparation for IV infusion**

#### **Labelling of prepared solution**

The label of the resulting solution must contain the following statement: THIS SOLUTION CONTAINS SORBITOL.

### **Reconstituting the Powder**

Appropriate aseptic technique should be used.

OGIVRI should be carefully handled during reconstitution. Causing excessive foaming during reconstitution or shaking the reconstituted OGIVRI may result in problems with the amount of OGIVRI that can be withdrawn from the vial.

Each 150 mg vial should be reconstituted with 7.2 mL of sterile water for injections as the solvent. The use of other solvents should be avoided. The resultant solution is 7.4 mL of approximately 21 mg/mL trastuzumab. A 4% overage is included to ensure withdrawal of the labelled dose of 150 mg.

Each 440 mg vial should be reconstituted with 20 mL of bacteriostatic water for injection as the solvent. The use of other solvents should be avoided. The resultant multiple-dose solution contains approximately 21 mg/mL trastuzumab. In patients with known hypersensitivity to benzyl alcohol, reconstitute with 20 mL of Sterile Water For Injection (SWFI) without preservative to yield a single use solution.

### **Instructions for Reconstitution**

150 mg vial



1. Using a sterile syringe, slowly inject 7.2 mL (for 150 mg vial) of sterile water for injections in the vial containing the lyophilized OGIVRI, directing the stream into the lyophilized cake.
2. Swirl vial gently to aid reconstitution. OGIVRI may be sensitive to shear-induced stress, e.g. agitation or rapid expulsion from a syringe. DO NOT SHAKE.

Slight foaming of the product upon reconstitution is not unusual. Allow the vial to stand undisturbed for approximately 5 minutes. The reconstituted preparation results in a colourless to pale yellow transparent solution and should be essentially free of visible particulates.

440 mg vial with bacteriostatic water for injection

1. Using a sterile syringe, slowly inject the 20 mL of diluent into the vial containing the lyophilized cake of OGIVRI. The stream of diluent should be directed into the lyophilized cake.
2. Swirl the vial gently to aid reconstitution. DO NOT SHAKE.
3. Slight foaming of the product may be present upon reconstitution. Allow the vial to stand undisturbed for approximately 5 minutes.
4. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Inspect visually for particulates and discoloration. The solution should be free of visible particulates, clear to slightly opalescent and colorless to pale yellow.
5. Store reconstituted OGIVRI at 2°C to 8°C; discard unused OGIVRI after 24 hours.

Note: If OGIVRI is reconstituted with SWFI without preservative (for patients with hypersensitivity to benzyl alcohol), use immediately and discard any unused portion.

**Instructions for Dilution**

*Weekly Regimen:* Determine the volume of the reconstituted solution required based on a loading dose of trastuzumab 4 mg/kg body weight, or a maintenance dose of trastuzumab 2 mg/kg body weight:

$$\text{Volume (mL)} = \frac{\text{Body weight (kg)} \times \text{dose (4 mg/kg for loading or 2 mg/kg for maintenance)}}{21 \text{ (mg/mL, concentration of reconstituted solution)}}$$

*Three-Weekly Regimen:* Determine the volume of the reconstituted solution required based on a loading dose of trastuzumab 8 mg/kg body weight, or subsequent every 3 weeks dose of 6 mg/kg body weight:

$$\text{Volume (mL)} = \frac{\text{Body weight (kg)} \times \text{dose (8 mg/kg for loading or 6 mg/kg for maintenance)}}{21 \text{ (mg/mL, concentration of reconstituted solution)}}$$

**Preparation and Stability of the Admixture**

The appropriate amount of the reconstituted solution should be withdrawn from the vial and added to an infusion bag containing 250 mL of 0.9% sodium chloride.

Dextrose (5%) solution should not be used since it causes aggregation of the protein. OGIVRI SHOULD NOT BE MIXED OR DILUTED WITH OTHER MEDICINES. No incompatibilities between OGIVRI and polyvinylchloride, polyethylene or polypropylene bags have been observed.

The infusion bag should be gently inverted to mix the solution in order to avoid foaming. Parenteral drug products should be inspected visually for particulates and discoloration prior to administration.

From a microbiological point of view, the OGIVRI infusion solution, if reconstituted with sterile water (SWFI) without preservative, should be used immediately. If diluted aseptically, it may be stored for 24 hours when refrigerated at 2°C to 8°C.

## OVERDOSAGE

There is no experience with overdosage in human clinical trials. Single doses higher than 10 mg/kg have not been tested.

Treatment of overdose should consist of general supportive measures.

For information on the management of overdose, contact the Poisons Information Centre (in Australia call 13 11 26; in New Zealand call 0800 767 766).

## PRESENTATION AND STORAGE CONDITIONS

OGIVRI powder for IV infusion is available as:

- single-dose 150 mg vial
- 440 mg vial with bacteriostatic water for injection

The 440 mg vial with BWFI is a Pharmacy Bulk Pack.

The contents of the sterile vial appear as a lyophilized, white to pale yellow powder. The reconstituted OGIVRI solution contains approximately 21 mg/mL of trastuzumab.

### Excipient with known effect

#### Sorbitol

Ogivri contains 0.77 mg sorbitol/ 1.00 mg trastuzumab. There is 115.2 mg of sorbitol in the 150 mg vial and 337.9 mg of sorbitol in the 440 mg Pharmacy Bulk Pack.

### Storage

Store OGIVRI 150 mg and 440 mg vials at 2 to 8°C. Refrigerate. Do not freeze. Do not use beyond the expiration date stamped on the vial.

### Reconstituted Solution

#### 150 mg

A vial of OGIVRI reconstituted with sterile Water for Injections (SWFI) without preservative should be used immediately and any unused portion must be discarded. OGIVRI 150 mg is for single use in one patient only. Do not freeze the reconstituted solution.

#### 440 mg

When reconstituted with BWFI, USP (1.1% benzyl alcohol) the Pharmacy Bulk Pack solution may be kept for no longer than 24 hours when stored at 2°C to 8°C or 6 hours at room temperature or above

### **Diluted Solution for Infusion**

From a microbiological point of view, the OGIVRI infusion solution, when reconstituted with sterile water (SWFI) without preservative, should be diluted and used immediately. The product is not intended to be stored after dilution.

Solutions of OGIVRI for infusion are physically and chemically stable in polyvinylchloride, polyethylene or polypropylene bags containing 0.9% sodium chloride at 2°C to 8°C for 24 hours, or 6 hours at room temperature or above.

### **Disposal of Medicines**

The release of medicines into the environment should be minimized. Medicines should not be disposed of via wastewater and disposal through household waste should be avoided. Unused or expired medicine should be returned to a pharmacy for disposal.

## **NAME AND ADDRESS OF THE SPONSOR**

### **Alphapharm Pty Limited**

Level 1, 30 The Bond

30-34 Hickson Road

Millers Point NSW 2000

ABN 93 002 359 739

[www.mylan.com.au](http://www.mylan.com.au)

## **POISON SCHEDULE OF THE MEDICINE**

S4 (Prescription Only Medicine)

## **DATE OF FIRST INCLUSION IN THE AUSTRALIAN REGISTER OF THERAPEUTIC GOODS (THE ARTG)**

11/12/2018

OGIVRI\_pi\Dec18/01