<u>17377</u>

Package leaflet: Information for the patient



Tiotropium (as bromide) 13 micrograms

Powder for inhalation (in Capsule)



# Braitus Tiotropium (as bromide)

#### Powder for inhalation

- This leaflet answers some common questions about Braltus Braltus capsules are supplied with a Zonda inhaler.
- It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.
- All medicines have benefits and risks. Your doctor has weighed the risks of you taking Braltus against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet.

You may need to read it again.

# What Braltus is used for?

Braltus is used to make breathing easier for people with chronic obstructive pulmonary disease (COPD).

This helps to improve your condition and to prevent exacerbations (periodic worsening of symptoms) from occurring.

Braltus improves breathing by relaxing the air passages that carry air to and from the lungs. It begins to act within 30 minutes after use

and the effect should last a full day. Braltus contains the active ingredient tiotropium (as bromide). It belongs to a group of medicines called anticholinergics.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

#### Before you take Braitus

When you must not take it Do not take Braltus if you have an allergy to:

- any medicine containing tjotropium bromide
- any of the ingredients listed at the end of this leaflet
- any medicine containing atropine or its derivatives, e.g. ipratropium or oxitropium
- any other anticholinergic medicines

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

Do not take this medicine to treat a sudden attack of breathlessness, wheezing or coughing.

You will need a different type of medicine.

Do not give this medicine to children or adolescents (below the age of 18 years).

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering. If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it
Tell your doctor if you have
allergies to any other medicines,
foods, preservatives or dyes.
Tell your doctor if you have or
have had any of the following
medical conditions:

- high pressure in the eye (glaucoma)
- kidney or liver problems
- problems with your prostate gland
- problems with passing urine
- you have suffered from a heart attack during the last 6 months or from any unstable or life threatening irregular heart beat or severe heart failure within the past year

Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.

Your doctor can discuss with you the risks and benefits involved. Braltus is not generally recommended for use in pregnant women.

If you have not told your doctor about any of the above, tell him/her before you start taking Braltus. Do not allow the powder to enter into the eyes. Should this occur, immediately flush your eyes with cold tap water for several minutes and immediately consult your doctor for further advice. If the powder enters the eye, it may result in eye pain or discomfort, blurred vision, seeing halos around lights or coloured images in association with red eyes (i.e. narrow angle glaucoma).

Take this medicine only with the Zonda® device.

#### Taking other medicines

Tell your doctor or pharmacist if you are using any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Braltus may interfere with each other. These include other anticholinergic medicines used to treat COPD such as glycopyrronium, aclidiniuim, umeclidinium or ipratropium.

These medicines may be affected by Braltus or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

#### How to take Braitus

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet if you do not understand the instructions on the box, ask your doctor or pharmacist for help.

#### How much to take

The recommended dose for adults is 1 capsule to be inhaled, once a day.

One capsule provides the required daily dose of tiotropium (a delivered dose of 10 micrograms of tiotropium); do not tke more than the recommended dose.

How it is given Inhale the powder in the capsules only using the Zonda device.

Do not place a capsule directly into the mouthpiece of the device.

Do not swallow the capsules. Do not open the capsules.

If you open the capsules, the small amount of powder inside may be lost, or you may accidentally get the powder in your eyes.

Use the capsules within 60 days of opening the bottle.

Read the Directions for Use at the end of this leaflet for instructions on how to use Braltus capsules with the Zonda device.

Follow the Directions for Use carefully. Pierce the capsule only once using the Zonda device.

Do not allow the powder to enter into the eyes. Should this occur, immediately flush your eyes with cold tap water for several minutes.

If you have any problems inhaling Braltus capsules using the Zonda device ask your doctor or pharmacist for advice.

When to take it Take your medicine at about the same time each day.

Taking it at the same time each day will have the best effect. It will also help you remember when to take it

How long to take it Continue taking your medicine for as long as your doctor tells you.

This medicine helps to control your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

#### If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect

If you are not sure what to do, ask your doctor or pharmacist. If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (Australia 13 11 26) for advice, or go to Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Braltus. Do this even if there are no signs of discomfort or poisoning.
You may need urgent medical attention.

Symptoms of an overdose may include fast or irregular heart beat, blurred vision, nausea, stomach pain, dry mouth, constipation and difficulty passing urine.

# While you are taking Braltus

Things you must do
If you are about to be started on
any new medicine, remind your
doctor and pharmacist that you
are taking Braltus.

Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately.
Keep all of your doctor's appointments so that your progress can be checked.

Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

It is very important to follow the Directions for Use carefully. If your breathing becomes more difficult while you are taking this medicine, tell your doctor immediately. If you are a smoker, your doctor or pharmacist can advise you on the steps to take to quit smoking.

Things you must not do
Do not take Braltus to treat any
other complaints unless your
doctor tells you to.
Do not give your medicine to
anyone else, even if they have
the same condition as you.
Do not take Braltus more
frequently than once daily

Things to be careful of Be careful driving or operating machinery until you know how Braltus affects you.
This medicine may cause dizziness or blurred vision in some people. If you have any of these

symptoms, do not drive, operate machinery, or do anything else that could be dangerous.

#### Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Braltus. This medicine helps most people with COPD, but it can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- dry mouth: this is usually mild sore mouth, gums, or throat; swollen, red, sore tongue
- oral thrush
- hoarse voice
- nose bleeds
- cough
- dizziness
- constipation
- trouble sleeping
- sinusitis, a feeling of tension or fullness in the nose, cheeks and behind your eyes, sometimes with a throbbing ache, fever, stuffy nose and loss of the sense of smell.

These are mild side effects of the medicine, and usually short-lived.

# Tell your doctor as soon as possible if you notice any of the following:

- blurred vision
- seeing halos around lights or coloured images in association with red eyes; high pressure in the eye (glaucoma)
- · difficulty in swallowing
- heartburn
- difficulty in passing urine
- pain while passing urine, urinary tract infection, increased need and frequency in passing urine
   a worsening of breathing
- a worsening of breathing problems (induced by the inhalation process). Inhaled medicines such as Braltus may cause tightness of the chest, coughing, wheezing or breathlessness immediately after inhalation.

The above list includes serious side effects that may require medical attention.

#### If any of the following happen, tell your doctor immediately or go to Emergency at your nearest hospital:

severe pain in the stomach with bloating, gut cramps and vomiting
changes in heart rate (fast, slow or irregular), palpitations
allergic reaction (shortness of breath, wheezing or difficulty

breathing; swelling of the face, lips, tongue or other parts of the body, rash, itching or hives on the skin).

The above list includes very serious side effects. You may need urgent medical attention or hospitalization.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.
Other side effects not listed above may also occur in some people.

#### After taking Braltus

Cleaning

If necessary, you may wipe the mouthpiece of your Zonda inhaler after use with a dry cloth or tissue.

Discard the Zonda device after 30 uses. Do not reuse the inhaler for another bottle of capsules.

Storage

Keep your capsules in the bottle until it is time to take them. If you take the capsules out of the bottle they may not keep well. Keep your capsules in a cool, dry place where the temperature stays below 25°C. Do not store your capsules in the refrigerator or freezer. Use the capsules within 60 days of opening the bottle.

Do not store Braltus or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car. Heat and dampness can destroy some medicines.

Keen it where children cannot

Keep it where children cannot reach it.

A locked cupboard at least oneand-a-half metres above the ground is a good place to store medicines.

#### Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

#### **Product Description**

#### What it looks like

Braltus capsules are colourless and translucent. Braltus capsules contain a white powder. The Zonda inhaler has a green body and a cap with a white push button.

Braltus capsules contain only a small amount of powder, which means that the capsule is only partially filled. The amount of powder in each capsule is equivalent in size to the tip of a matchstick.

Braltus is available in a carton containing a bottle of 30 capsules with the Zonda device.

Braltus and Spiriva both deliver 10 micrograms of tiotropium and are equivalent.

#### Ingredients

Braltus capsules contains 13
micrograms of tiotropium
(equivalent to tiotropium bromide
15.6 micrograms) as the active
ingredient.
During inhalation, 10 micrograms
of tiotropium are delivered from
each capsule from the mouthpiece
of the Zonda inhaler.
It also contains lactose
monohydrate (which contains milk
protein) and Vcaps The Vegetarian
Alternative empty hard capsule
size 3 capsule (Ingredient ID
10127).

#### Sponsor

Braltus is supplied in Australia by: Teva Pharma Australia Pty Ltd 37 Epping Road Macquaire Park NSW 2113 Braltus<sup>®</sup> is a registered trademark of Teva Pharma

#### Zonda Instructions for Use

The Zonda inhaler enables you to inhale the medicine contained in the Braltus capsule that your doctor has prescribed for your breathing problems.

Remember to carefully follow your doctor's instructions for using Braltus. The Zonda inhaler is especially designed for Braltus capsules: you must not use it to take any other medication. The capsules must only be inhaled using the Zonda inhaler. Do not use any other inhalers to take Braltus capsules. Each capsule contains only a small amount of powder. Do not open the capsule or it may not work.

The Zonda inhaler should only be used with the bottle of capsules provided. Do not reuse the inhaler for another bottle of capsules. Discard the Zonda device after 30 uses.



- 1. Dust cap
- 2. Mouthpiece
- 3. Base
- 4. Piercing button
- 5. Centre chamber








- 1. Pull the cap upwards
- Hold the base of the inhaler firmly and open the mouthpiece by pulling it upwards, in the direction of the arrow.
- Remove a Braltus capsule from the bottle immediately before use and close the bottle tightly. Place one capsule in the capsule-shaped compartment in the base of the inhaler. Do not store the capsule in the Zonda inhaler.
- Never place a capsule directly into the mouthpiece.








- 5. Close the mouthpiece until you hear a click, leaving the cap open.
- 6. Hold the inhaler with the mouthpiece upwards, and firmly press the piercing button completely in only once. Release the button. This will pierce the capsule and allows the medication to be released when you breathe in.




- Breathe out fully. It is important to do this away from the mouthpiece. Avoid breathing into the mouthpiece at any time.
- 8. Place the mouthpiece in your mouth and keep your head in an upright position. Close your lips around the mouthpiece and breathe in slowly and deeply enough to hear or feel the capsule vibrating inside the compartment. Hold your breathe for as long as you comfortably can whilst taking the inhaler out of your mouth. Then breathe normally. Repeat steps 7 and 8 to empty the capsule completely.
- 9. After use, open the mouthpiece again, and tip out the empty capsule. Close the mouthpiece and cap, and store your Zonda inhaler.

#### Manufacturer:

Plastiape S.p.A. Via 1 Maggio, 8, 23875 Osnago LC, Italy





⊕Brattus Tiotropium (as bromide)

- Powder for inhalation

  This leaflet answers some common questions about Braitius Braitius Braitius Braitius Braitius Braitius Braitius Capaules are supplied with a Zonda inhaler.

  It does not contain all the available information. It does not take the place of taking to your doctor of pharmacists and risks, Your doctor has weighted the risks of your laking Braitius against the benefits they expect it will have for you taking Braitius against the benefits they expect it will have for your laking Braitius.

If you have any concerns about taking this medicine, askyour doctor or pharmacist.

Keep this leaflet.

You may need to read it again.

# What Braitus is used for?

Final Braitos is used for?

Braitus is used to make breathing caster for people with chronic obstructive pulmonary disease (COPD).

This helps to improve your condition and to prevent exactbations (periodic worsening of symptoms) from occurring.

Braitus improves breathing by relaxing the air passages that carry air to and from the lungs. It begins to act within 30 minutes after use and the effect should last a full day.

Braitus contains the active ingredient inclinging a group of medicines called anticholinergics.

Ask your doctor if you have any questions about why this medicine has been prescribed for you Your doctor may have prescribed if for another reason.

#### Before you take Brattus

Before you take Braitius

When you must not take it
Do not take Braitius if you have an
allengy to:

any medicine containing
totropium bromide
any of the lingredients
listed at the end of this
loaflet
any medicine containing
atropine or its derivatives,
e.g. ipratropium or
oxtropium
any other anticholinergic
medicines
Some of the symptoms of an
alkergic reaction may include
shortness of breath
wheeping or difficutly
to walking of the face, lips,
tongue or other parts of the
body
take hickning or hives on the
skin
Do not take this medicine to treat
a sudden attack of
breathlessness, wheezing or
coughing,
you will need a different type of
medicine
Do not give this medicine to
the action of the packaging is torn
or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal if you are not sure whether you should start taking this medicine, talk to your doctor.

snouls start taking ins medicine, talk to your doctor. Before you start to take it Yell your doctor if you have allorgies to any other medicines, foods, preservatives or dyes. Tell your doctor if you have or have had any of the following medical conditions:

Ingh pressure in the eye (glaucoma)

Kidney or liver problems

problems with your prostate gland

problems with passing unne
you have suffered from a heart attack during the last of months or from any unstable or like threatening regular heart beat or severe heart failure within the past year

Tell your doctor if you are pregnant or are breast-feeding. Your doctor can discuss with you the risks and benefits involved Brakus is not generally recommended for use in pregnant or about any of the above, tell hirnher before you start taking Braltus.

Do not allow the powder to enter Into the eyes, Should this occur, Immediately flush your eyes with cold tap water for several minutes and Immediately consult your doctor for further advice. If the powder enters the eye, it may result in eye pain or discomfort, burred vation, seeing halos around lights or coloured mages in association with red eyes (i.e. narrow angle glaucoma).

Take this medicine only with the Zondas device.

Taking other medicines

Tell your doctor or pharmacist if you are using any other modicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Braitus may atterfere with each other These include other anticholinergic medicines used to treat COPD such as glycopyrsomm, actidinium, umecidinum or ipratropium.

These medicines may be affected by Braitus or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while laking this medicine

#### How to take Braitus

Follow all directions given to you by your doctor or pharmacist carefully They may differ from the information contained in this leaflet. If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

Instructions on market and the pode of the pharmacist for help.

How much to take The recommended dose for adults is 1 capsule to be inhaled, once a day One capsule provides the required daily dose of fottopsium (a delivered dose of 10 micrograms of biotopsium), do not tike more than the recommended dose How it is given inhale the powder in the capsules only using the Zonda device.

Do not place a capsule directly into the mouthpiece of the device.

Do not place a capsule s. If you open the capsules.

If you open the capsules.

If you open the capsules may be lost, or you may accidentally get the powder in your eyes.

Use the capsules within 60 days

Use the capsules within 60 days of opening the bottle.

Read the Directions for Use at the end of this leaflet for Instructions on how to use Brallus capsules with the Zonda device.

Follow the Directions for Use carefully. Pierce the capsule only once using the Zonda device.

Do not allow the powder to enter into the eyes. Should this occur, immediately flush your eyes with cold tap water for several minutes. If you have any problems inhaling Braitus capsules using the Zonda device ask your doctor or pharmacist for advice.

When to take it Take your medicine at about the same time each day.

Taking it at the same time each day will have the bast effect. It will also help you remember when to take it. How long to take it. Continue taking your medicine for as long as your doctor tells you.

for as tong as you. According you.
This medicine helps to control your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

If you forget to take it

If you forget to take it

If it is almost time for your next
dose, skip the dose you missed
and take your next dose when
you are meant to.
Otherwise, take it as soon as you
remember, and then go back to
taking your medicine as you
would normally.
Do not take a double dose to
make up for the dose that you
missed.
This may increase the chance of
you getting an unwanted side
effect.
If you have trouble remembering
to take your medicine, ask your
pharmacist for some hints.
If you take too much feverfose!

If you take too much (overdose)

If you take too much (everdose) immediately telephone your doctor or the Poisons information Centire (Australia 13 11 26) for advice, or go to Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Braitus. Do this even if there are no signs of discomfort by poisoning. You may need urgent medical attention.

Symptoms of an overdose may include fast or irregular heart beat, blurred vision, nausea, stomach pain, dry mouth, constipation and difficulty passing urine.

## While you are taking Braitus

Braitus

Things you must do
If you are about to be started on
any new medicine, remind your
doctor and pharmacist that you
are taking Braitus.
Tell any other doctors, dentists,
and pharmacists who treat you
that you are taking this
medicine.
If you are going to have surgery
tell the surgeon or anaesthelist
that you are taking this
medicine.
It may affect other medicines used
during surgery
If you become pregnant while
taking this medicine, tell your
doctor immediately.
Keep all of your doctor's
appointments so that your
progress can be checked.
Your doctor may do some tests

Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

It is very important to follow the Directions for Use carefully. If your breathing becomes more difficult while you are taking this medicine, tell your doctor immediately. If you are a smoker, your doctor or pharmacist can advise you on the steps to take to quit smoking.

Things you must not do
Do not take Braitus to treat any
other complaints unless your
doctor tells you to.
Do not give your medicine to
anyone else, even if they have
the same condition as you.
Do not take Braitus more
frequently than once dally

Things to be careful of Be careful driving or operating machinery until you know how Braitus affects you. This medicine may cause dizziness or blurred vision in some people if you have any of these symptoms, do not drive, operate machinery, or do anything site that could be dangerous.

#### Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Braitus.

This medicine helps most people with COPD, but it can have side effects. Sometimes they are entous, most of the time they are not. You may need medical attention if you got some of the side effects. You may not experience any of them.

Ask your dector or pharmacist to answer any questions you may have.
Tell your dector or pharmacist to answer any questions you may have.
Tell your dector or pharmacist if you notice any of the following and they worry you:
Any mouth this is usually mild sore mouth, gums, or threat, swellen, red, sore tongue oral thrush hoarse voce nose bleeds cough dizziness consipation trouble sleeping sinusitis, a feeling of tension or fullness in the nose, cheeks and behind your yees, sometimes whe a throbbing ache, fever, stuffy nose and loss of the sense of smell.

Tell your doctor as eoon as possible if you notice any of the following:

- blurred vision
- seeing halos around lights or coloured images in association with red eyes, high pressure in the eye (glaucoma)
- difficulty in wallowing
- heartburn
- pain while passing urine, urinary tract infection, increased need and frequency in passing urine, urinary tract infection, increased need and frequency in passing urine, a worstening of breathing problems (induced by the inhalation process) Inhaled modicines such as Brahus may cause lightness of line chest coughing, wheeching or breathlessness immediately after inhalation
- The above list includes serious side effects that may require medical attention.

If any of the following happen,

medical attention.

If any of the following happen, tell your doctor immediately or go to Emergency at your nearest hospital:
- severe pain in the stomach with booling, gut cramps and vomiting changes in heart rate (fast, slow or irregular), palgitations.
- allergic reaction (shortness of breath, wheezing or difficulty

breathing, swelling of the face, tips, longue or other parts of the body, rash, tiching or hives on the skinly. The above list includes very serious side effects. You may need urgent medical attention or hospitatization. Tell your doctor or pharmaclet if you notice anything that is making you feel unwell. Other side effects not listed above may also occur in some people.

### After taking Braitus

Cleaning
If necessary, you may wipe the mouthpiece of your Zonda inhaler after use with a dry cloth or tissue.

or tissue.

Discard the Zonda device after 30 uses. Do not reuse the inhalor for another bottle of capsules.

Storage
Keep your capsules in the bottle until it is time to take them. If you take the capsules out of the bottle they may not keep well keep your capsules in a cool, dry place when the temperature stays below 25°C. Do not store your capsules in the refrigerator or freeze. Use the capsules within 60 days of opening the bottle.

Do not store Braitus or any other medicine in the bathroom or near a sink. Do not leave it on window all or in the car. Heat and dampness can destroy some medicines
Keep it where children cannot reach it.
A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines

Disposal
If your doctor tells you to stop
taking this medicine or the
explry date has passed, ask
your pharmacist what to do with
any medicine that is left over.

#### Product Description

What it looks like Bratus capsules are colourless and translucent Bratus capsules are colourless and translucent Bratus capsules contain a write provider. The Zonda inhaler has a green body and a cap with a write push button.

Bratus capsules contain only a small amount of powder, which means that the capsule is only partially filled. The amount of powder in each capsule is only settled to a matchastick.

Bratus is available in a carton containing a bottle of 30 capsules with the Zonda device.

Braltus and Spiriva both deliver 10 micrograms of flotropium and arc bagulvalent. Ingredlents Braltus capsules contains 13 micrograms of liotropium equivalent to lotropium bromide 15 6 micrograms of liotropium gredlents. During inhalation, 10 micrograms of lotropium are delivered from each capsule from the mouthplece of the 2 ond as inhaler. It also contains lactose monohydrate (which contains mik protein) and Vcaps The Vegetarian Alternative empty hard apsule site 3 capsule (ingredient ID 10127).

Sponsor Bratus is supplied in Australia by Teve Pharma Australia Pty Ltd 37 Epping Road Macquarie Park NSW 2113 Bratus\* is a registered trademark of Teva Pharma

Commented (SLI): As per recommendation Quality report

#### Zonda Instructions for Use

The Zonda inhalor enables you to inhale the medicine contained in the Brattus capsule that your doctor has prescribed for your breathing problems

Remember to carefully follow your doctor's instructions for using finatus. The Zonda inhaler in especially designed for Braitus capsules; your must not use it take any other medication. The capsules must only be inhaled using the Zonda inhaler. Do not use any other inhalers to take Braitus capsules. Each capsule contains only a small amount of powder. Do not open the capsule or it may not work. The Zonda inhaler should only be used with the bottle of capsules provided. Do not reuse the inhaler for another bottle of capsules. Discard the Zonda device after 30 uses.









1 Pull the cap upwards
2 Hold the base of the inhaler firmly and open the mouthpiece by pulling a upwards, in the direction of the arrow.


Remove a Braitus capsule from the bottle immediately before use and close the bottle tightly Place one capsule in the capsule-shaped compartment in the base of the inhaler Do not store the capsule in the Zonda inhaler.

Never place a capsule directly into the mouthpiece until you hear a click, leaving the cap open.

Hold the inhaler with the mouthpiece upwards, and firmly piecs the piecing button completely in only once Release the button. This will piece the capsule and allows the medication to be released when you breathe in.







- 7 Breathe out fully. It is important to do this away from the mouthpiece. Avoid breathing into the mouthpiece at any time.
  8 Place the mouthpiece in your mouth and keep your head in an upright position. Close your place the capsule whating inside the compartment Hold your breathe for as long as you comfortably can whilst taking the makeler out of your mouth. Then breathe for as long as you comfortably can whilst taking the makeler out of your mouth. Then breathe normally Repeat steps 7 and 8 to empty the capsule completely. After use, open the mouthpiece again, and to out the empty gaspule. Close the mouthpiece and cap, and store your Zonda intaler.

  Manufacturer:

Manufacturer: Plastiape S p A Via 1 Maggio, 8, 23875 Osnago LC, Baly

